Preprint typeset in JHEP style - HYPER VERSION

CALT-68-2843

arXiv:1107.0973v1 [hep-th] 5 Jul 2011

Towards a 4d/2d correspondence for Sicilian quivers

Lotte Hollands[*], **Christoph A. Keller**[†], **Jaewon Song**[‡]

California Institute of Technology, Pasadena, CA 91125, USA

Abstract: We study the 4d/2d AGT correspondence between four-dimensional instanton counting and two-dimensional conformal blocks for generalized $SU(2)$ quiver gauge theories coming from punctured Gaiotto curves of arbitrary genus. We propose a conformal block description that corresponds to the elementary $SU(2)$ trifundamental half-hypermultiplet, and check it against $Sp(1) - SO(4)$ instanton counting.

[*]hollands@caltech.edu

[†]ckeller@theory.caltech.edu

[‡]jaewon@theory.caltech.edu

Contents

1. Introduction **1**

2. Instanton counting for half-hypermultiplets **7**
- 2.1 Half-hypermultiplets 7
- 2.2 Instanton counting for half-hypermultiplets 9

3. CFT building blocks for Sicilian quivers **13**
- 3.1 Three-point functions 14
- 3.2 Partition function for the trifundamental coupling 16

4. Towards a 4d/2d correspondence for Sicilian quivers **18**

5. Examples **21**
- 5.1 The $SU(2)$ trifundamental as a $Sp(1) - SO(4)$ bifundamental 22
- 5.2 Genus two quiver through $Sp(1) - SO(4)$ instanton counting 25
- 5.3 Alternative prescriptions for the genus two quiver 29

6. Discussion **31**

A. More about the trifundamental half-hypermultiplet **33**

B. Contour integrands for Sicilian quivers **36**

1. Introduction

The Alday-Gaiotto-Tachikawa correspondence [1] relates the instanton partition functions of $\mathcal{N} = 2$ gauge theories to chiral blocks. Such theories can be engineered in M-theory by wrapping M5-branes on a punctured Riemann surface $\mathcal{C}$, the so-called Gaiotto curve [2]. The correspondence is obtained by decomposing $\mathcal{C}$ into pairs of pants, and computing the corresponding conformal block.

If the corresponding Gaiotto curve $\mathcal{C}$ is of genus zero or one, a weakly coupled description exists in certain regions of its complex structure moduli space. The gauge theory is then described by a linear or cyclic quiver, and it corresponds to a linear or cyclic decomposition of the Gaiotto curve. For these cases the correspondence between conformal blocks and instanton partition functions has been checked extensively in the literature [1, 3, 4].

For Sicilian gauge theories of A_n-type with $n \geq 2$, however, such a Lagrangian description does not exist in general. When decomposing $\mathcal{C}$ into pairs of pants we encounter various strongly interacting isolated four-dimensional SCFT's whose flavor symmetries are partially gauged [2, 5], such as the ones that appear in the Argyres-Seiberg duality [6].

For A_1 theories the situation is better: Sicilian gauge theories of A_1-type admit a Lagrangian description throughout the complex structure moduli space of the Gaiotto curve $\mathcal{C}$. In this case, a pair of pants just refers to a trifundamental half-hypermultiplet in the four-dimensional gauge theory. Since instanton counting is formally developed for any $\mathcal{N} = 2$ gauge theory with a Lagrangian prescription, it should be possible to write down instanton partition functions for Sicilian $SU(2)$ quiver gauge theories. Moreover on the CFT side it should then be possible to write down conformal blocks corresponding to such decompositions. Having done this, we should be able to compare the two and check if the AGT correspondence still works in this case. To our knowledge this has not yet been accomplished in the literature, and only indirect checks in this direction have been performed [7].

Unfortunately, this extension to Sicilian quivers is not that straightforward, and introduces a number of subtleties on both sides of the story.

First of all, note that the conventional method to count instantons in $SU(2)$ gauge theories is to consider instanton counting for gauge group $U(2)$, and impose the tracelessness condition at the end to reduce to $SU(2)$. This $U(2)$ prescription follows from resolving the ultra-violet singularities of the instanton moduli space by turning on an FI parameter. Mathematically, it can be elegantly formulated in terms of counting rank two torsion-free sheaves.

It is important however to realize that Sicilian quivers are in general *not* defined for gauge group $U(2)$. The reason for this is that the trifundamental field, that couples three $SU(2)$ gauge groups, is described by a half-hypermultiplet. (A full trifundamental hypermultiplet contains too many degrees of freedom, so that the resulting theory would not be conformal. [1]) Gauge theories involving half-hypermultiplets are only CPT invariant if the corresponding matter fields transform in a pseudo-real representation. (We will present a detailed explanation of this in section 2.) The fundamental representation of $SU(2)$ is indeed pseudo-real, whereas the fundamental of $U(2)$ is complex. It is therefore not possible to just work with the $U(2)$ as before and specialize to the $SU(2)$ case in the end.

On the face of it there is one obvious way around this problem. Instanton counting for $Sp(N)$ and $SO(N)$ gauge groups has been developed in [8], and the D-type quivers relevant in this context were investigated in [9]. [2] We can thus try to use the fact that $Sp(1) = SU(2)$ to circumvent those issues and compute the $SU(2)$ instanton parition function directly. Similarly, we can also use $SO(4) = SU(2) \times SU(2)$ to compute configurations involving bifundamentals

[1] One exception to this is the genus two quiver, which describes three $SU(2)$ gauge groups coupled by two $SU(2)$ trifundamental half-hypermultiplets, and can be equivalently described by a full $U(2)$ trifundamental hypermultiplet. We will come back to this example in section 5.

[2] See also [10] for a discussion of such theories.

of $SU(2)$.

This approach was explored in [4]. Rather surprisingly it was found that the $Sp(1)$ and the $U(2)$ instanton counting schemes yield seemingly different results for conformal quiver gauge theories, even though by the above remarks they should describe the same physics. The resolution of this apparent contradiction is that the instanton partition functions are a priori expressed in terms of microscopic coupling constants. To compare the results, they need to be expressed first in terms of the physical IR coupling, after which they indeed agree. To put it another way, the results in the UV are related by a non-trivial mapping of microscopic coupling constants, and thus by a different choice of renormalization scheme for the conformal gauge theory. For theories that have a string theory embedding, this mapping has an elegant geometric interpretation in terms of the corresponding Gaiotto curves. [3]

Let us briefly recapitulate one of the examples of [4], the conformal $SU(2)$ gauge theory coupled to four hypermultiplets. The $Sp(1)$ and $U(2)$ instanton partition functions are related by the mapping [4]

$$q_{U(2)} = q_{Sp(1)} \left(1 + \frac{q_{Sp(1)}}{4}\right)^{-2}, \tag{1.1}$$

where $q_{U(2)}$ is the microscopic coupling for the $U(2)$ scheme and $q_{Sp(1)}$ that of the $Sp(1)$ scheme. Geometrically, this mapping identifies the cross-ratios of the corresponding $U(2)$ and $Sp(1)$ Gaiotto curves, which are related to each other by a double cover construction. This is illustrated in figure 1.

Another way of phrasing all this is that the Nekrasov-Shadchin method of instanton counting singles out a particular choice of coordinates on the complex structure moduli space of the Gaiotto curve. For the AGT correspondence to work, it is clear that we need to take the same coordinates on the moduli space of the conformal block as well. For standard quiver theories, the $U(2)$ parametrization of the moduli space corresponds to the standard choice of CFT coordinates for the punctured sphere and torus. The two sides thus agree immediately. For linear and cyclic Sp/SO quivers, the choice of coordinates is slightly more involved [4], but has a very natural interpretation too. For Sicilian quivers it is no longer obvious what coordinates to pick, and in fact most choices have unappealing features, as we will discuss below.

Let us emphasize that these complications only arise for conformal quiver gauge theories, and not for asymptotically free ones. For asymptotically free theories, one can *not* have a non-trivial mapping of parameters because the instanton expansion parameter $q = \Lambda^{b_0}$ is dimensionful, where b_0 is the coefficient of the beta function. So the instanton counting for all

[3] Geometrically, the physical IR coupling corresponds to the period matrix of the Seiberg-Witten curve, which is a branched covering over the Gaiotto curve. Inequivalent coverings lead to non-trivial mappings of the microscopic UV coupling constants. Mathematically, such a UV-UV mapping defines an isomorphism between Hitchin systems with the same spectral curve.

[4] We have checked this up to order 6 in the microscopic couplings.



Figure 1: The double covering of the the $U(2)$ Gaiotto curve over the $Sp(1)$ Gaiotto curve relates the microscopic coupling $q_{U(2)}$ (which is the cross-ratio of the $U(2)$ Gaiotto curve) to the microscopic coupling $q_{Sp(1)} = 4q$. Note that, whereas a $U(2)$ gauge group is represented by a tube, an $Sp(1)$ gauge group is represented geometrically by a tube with a twist-line. In the double covering this twist-line gets the interpretation of a branch-cut.

possible realizations of an asymptotically free gauge theory (such as for a single trifundamental field coupled to three $SU(2)$ gauge groups) should agree directly. On the CFT side this means that the corresponding correlation function should be essentially independent of the choice of parametrization.

After these remarks, let us turn to our main object of interest. As we have pointed out, the trifundamental building block of Sicilian $SU(2)$ quivers can be either described as a $Sp(1)$ trifundamental coupling or as a $Sp(1) - SO(4)$ bifundamental coupling. Physically, both of these couplings are equivalent to an $SU(2)$ trifundamental interaction. The $Sp(1) - SO(4)$ description has the advantage that it admits a type IIA string embedding using D4–NS5 branes and O-branes. Yet it only parametrizes a subspace of the moduli space of the $SU(2)$ trifundamental, as it involves just two gauge couplings instead of three. This is the exact opposite of the $Sp(1)^3$ description, which cannot be embedded in string theory, but does parametrize the whole gauge coupling moduli space. This is illustrated in figure 2.

Whether or not a particular realization of a given $\mathcal{N} = 2$ SCFT has a string embedding is important when we wish to find its dual 2d CFT description. After all, we expect to find a conformal field theory that lives on the corresponding Gaiotto curve.

Finding a conformal field theory prescription for the $SU(2)$ trifundamental half-hyper is conceptually easy. The general rule is that punctures coming from hypers correspond to insertions of primary fields, and punctures coming from the cutting of tubes correspond to insertions of descendant fields. The building block corresponding to a bifundamental is thus the correlator of one primary and two descendant fields. It is thus natural to expect that trifundamental building block should be a three-point function with three descendant fields



Figure 2: The Gaiotto curve for the $SU(2)$ trifundamental as a double cover over the $Sp(1) - SO(4)$ Gaiotto curve (on the left) and the non-existing $Sp(1)^3$ Gaiotto curve (on the right). On the right is illustrated why we cannot find a Gaiotto curve corresponding to the $Sp(1)$ trifundamental: it is not possible to close the three twist-lines (or branch-cuts) on the $Sp(1)$ curve.

inserted

$$\langle V(\phi^1_{I_1}, z_1) V(\phi^2_{I_2}, z_2) V(\phi^3_{I_3}, z_3) \rangle \ . \tag{1.2}$$

Since the half-hyper is massless and does not give a puncture in the Gaiotto curve, it makes sense not to insert a corresponding vertex operator. In our notation the field $\phi^i_{I_i}$ is a Virasoro descendant, and $V(\phi^i_{I_i}, z_I)$ is the vertex operator of the field $\phi^i_{I_i}$ inserted at position z_i. The weights of the fields $\phi^i_{I_i}$ encode the Coulomb branch parameter of an $SU(2)$ gauge group.

The main issue here is the choice of the insertion points z_i. For the standard choice of coordinates for linear decompositions they turn out as $0, 1, \infty$. For Sicilian decompositions it is no longer clear what the correct choice is. More precisely, different choices give different parametrizations of the moduli space. If we want to match conformal block to a given instanton counting result, we need to make sure to pick the correct prescription.

Alternatively, we can also be less ambitious and simply check that the results agree once we express them in terms of IR variables (*i.e.* as objects defined on the Seiberg-Witten curve). In particular we can circumvent this entire issue by considering asymptotically free theories, where the relation between UV and IR is trivial. In this way we can find the conformal block dual to the $SU(2)$ trifundamental half-hypermultiplet.

We can either start from the conformal quiver theory with six massive fundamental hypermultiplets and take a decoupling limit in which we send all the six masses to infinity. This limit decouples the hypermultiplets and leaves the three $SU(2)$ gauge groups coupled by a single trifundamental half-hypermultiplet, see figure 3. Alternatively, we can compute



Figure 3: On the left is illustrated the Gaiotto curve for the conformal quiver gauge theory with three $SU(2)$ gauge groups that are all coupled by a trifundamental interaction and each individually to two massive fundamental hypermultiplets. In the decoupling limit where we take all masses m_i of the fundamental hypermultiplets to infinity, we are left with the Gaiotto curve corresponding to the asymptotically free quiver gauge theory that couples the three $SU(2)$ gauge groups by a trifundamental half-hypermultiplet.

the conformal block as the correlation function of three Gaiotto states,

$$\boxed{\langle h_1, \Lambda_1 | V(|h_2, \Lambda_2\rangle, 1) | h_3, \Lambda_3 \rangle} \tag{1.3}$$

where the Gaiotto state $|h, \Lambda\rangle$ is an eigenstate of L_1 with eigenvalue Λ. The state $|h, \Lambda\rangle$ has appeared in the dual conformal field theory description of asymptotically free linear quiver theories, where it describes the asymptotic boundary conditions of the quadratic differential on the Gaiotto curve [11].

Finding the conformal block (1.3) dual to the $SU(2)$ half-trifundamental is one of the main results of this paper. We verify this prescription by checking several consistency requirements and by comparing it with instanton counting using the $Sp(1)-SO(4)$ scheme. We furthermore propose a prescription for the 4d/2d correspondence for any Sicilian quiver, and check this in several examples.

The outline of this paper is as follows: We start in section 2 with an introduction to half-hypermultiplets and particularly to instanton counting for half-hypermultiplets. The result of this exercise is a contour integral for the $Sp(1) - SO(4)$ bifundamental half-hypermultiplet. We continue in section 3 with the dual conformal field theory prescription and find the three-point function (1.3). In section 4 we formulate the 4d/2d correspondence between conformal blocks and generalized A_1 quivers. We check this proposal in several examples in section 5. Here we pay special attention to conformal quivers, where we find a non-trivial mapping of microscopic couplings. We conclude and discuss several directions for further research in section 6. Appendix A contains some more background on the $SU(2)$ trifundamental half-hyper, whereas appendix B summarizes the relevant contour integrands for Sicilian quivers.

2. Instanton counting for half-hypermultiplets

The trifundamental fields that appear in Sicilian quiver gauge theories of type A_1 form half-hypermultiplet representations of the $\mathcal{N} = 2$ SUSY algebra, in contrast to the more common (full) hypermultiplets. In this section we review the basic properties of Sicilian quivers with trifundamental half-hypermultiplets and show that they preserve $\mathcal{N} = 2$ supersymmetry. Subsequently, we develop the tools for counting instantons in quiver gauge theories with half-hypermultiplets. We apply these tools in section 5 to compute the instanton partition functions of Sicilian quivers.

2.1 Half-hypermultiplets

There are two types of $\mathcal{N} = 2$ supersymmetry multiplets: the $\mathcal{N} = 2$ vector multiplet and the hypermultiplet. The former consists of a vector field A_μ, two Weyl fermions λ_α and ψ_α, and one complex scalar B. All of them transform in the adjoint representation of the gauge group. In $\mathcal{N} = 1$ language such an $\mathcal{N} = 2$ vector multiplet consists of one $\mathcal{N} = 1$ vector multiplet with component fields (A_μ, λ_α) and one chiral multiplet with components (B, ψ_α). A hypermultiplet requires a choice of representation R of the gauge group. In $\mathcal{N} = 1$ language it consists of two chiral multiplets Q and $\tilde{Q}$, the former transforming in the representation R and the latter in its complex conjugate R^*. The chiral multiplet Q has component fields (ϕ, χ_α), and the anti-chiral multiplet $\tilde{Q}$ has components $(\tilde{\phi}, \tilde{\chi}_\alpha)$. We call both Q and $\tilde{Q}$ half-hypermultiplets.

The half-hypermultiplets Q and $\tilde{Q}$ form massless representations of the $\mathcal{N} = 2$ SUSY algebra. However, even though the helicities of the states in a half-hypermultiplet form a CPT complete distribution, the half-hypermultiplet does not transform as a real representation of the SUSY algebra. Indeed, notice that the massless $\mathcal{N} = 2$ SUSY algebra is equal to the Clifford algebra $\mathrm{Cl}_{4,0}$ with invariance group $SO(4)$. The four-dimensional representation of the Clifford algebra $\mathrm{Cl}_{4,0}$, under which the half-hypermultiplet transforms, is pseudo-real instead of (strictly) real. A generic half-hypermultiplet will therefore not be invariant under CPT.

It is possible though to circumvent this constraint. More precisely, an $\mathcal{N} = 2$ multiplet is CPT invariant if it transforms under a real representation of the product of the SUSY algebra, the gauge group and possible flavor groups. So, apart from the obvious possibility of combining a chiral and an anti-chiral multiplet into a full hypermultiplet, we can also consider a single half-hypermultiplet in a pseudo-real representation of the gauge group G.

Nevertheless, there is an additional requirement. Even when a single half-hyper-multiplet transforms in a real representation of $\mathrm{Cl}_{4,0} \times G$, such a theory may still be anomalous due to Witten's anomaly argument [12]. According to this argument for example a single half-hypermultiplet in the fundamental representation of $SU(2)$ is anomalous (its partition function vanishes) since it contains an odd number of chiral fermions. On the contrary, a single

half-hypermultiplet in the fundamental representation of $SU(2)^3$ contains four chiral fermions in each $SU(2)$-representation. Therefore, quiver gauge theories with $SU(2)$ trifundamental half-hypermultiplets are free of Witten's $SU(2)$ anomaly as well as CPT invariant.

Other examples of consistent theories with half-hypermultiplets occur when we consider massless bifundamental couplings between SO and Sp gauge groups. A half-hypermultiplet transforming under the bifundamental of $SO \times Sp$ is in a pseudo-real representation of the gauge group $G = SO \times Sp$ and is free of the Witten anomaly as well.

Half-hypermultiplet as a constrained hypermultiplet

Since working with full hypermultiplets is often much more efficient than with half-hypermultiplets, in what follows we find an alternative method to deal with half-hypers. Instead of considering a half-hypermultiplet by itself, we start with a full hypermultiplet (consisting of two half-hypermultiplets) and impose a constraint on it which only leaves a half-hypermultiplet.

A full hypermultiplet can be thought of as a multiplet formed out of two $\mathcal{N} = 1$ chiral multiplets Q and $\tilde{Q}$. The chiral multiplet Q transforms in representation R and the anti-chiral multiplet $\tilde{Q}$ in its complex conjugate R^*. By the remarks above, for the theory of a single half-hyper to make sense, R needs to be a pseudoreal. Let σ_G be the anti-linear involution that maps the representation R to its complex conjugate R^*. Since the representation R is pseudo-real, it obeys $\sigma_G^2 = -1$. For example, in the case of the fundamental representation of $SU(2)$, the involution σ_G is given by the ϵ-tensor $i\sigma_2$. For the trifundamental representation of $SU(2)$ it is given by the product of three ϵ-tensors, one for each $SU(2)$ gauge group.

To impose our constraint, we need a map τ that relates Q to $\tilde{Q}$ and vice versa. Since $\tilde{Q}$ appears in the complex conjugate, τ needs to be anti-linear. Moreover, it needs to preserve the representation, which means that it must involve σ_G. Let us write a full hypermultiplet as Q^a whose two half-hypermultiplet components are given by $Q^1 = Q$ and $Q^2 = \tilde{Q}^*$. The involution τ is then defined by

$$Q^a \mapsto \tau(Q^a) = \sigma_G \otimes \sigma_I (Q^a)^* \,, \tag{2.1}$$

where

$$\sigma_I \begin{pmatrix} Q^1 \\ Q^2 \end{pmatrix} = \begin{pmatrix} -Q^2 \\ Q^1 \end{pmatrix} . \tag{2.2}$$

It is straightforward to check that indeed $\tau^2 = 1$. We can describe a half-hypermultiplet as a hypermultiplet that stays (anti-)invariant under τ, *i.e.* that is an eigenvector of τ of eigenvalue ± 1. More explicitly, such a hyper is given by $(Q, \pm\sigma_G Q^*)$.

This description of a half-hypermultiplet is for instance convenient to find the Lagrangian for a half-hyper Q starting from the Lagrangian of a full hyper. Recall that the Lagrangian

for the hypermultiplet $Q^a = (Q, \tilde{Q}^*)$ coupled to a vector multiplet (V, Φ) is given by

$$\mathcal{L}_{\text{fh}} = \int d^2\theta\, d^2\bar{\theta} \left(Q^\dagger e^V Q + \tilde{Q}^t e^{-V} \tilde{Q}^* \right) + 2\sqrt{2}\ \text{Re} \int d^2\theta \left(\tilde{Q}^t\, \Phi\, Q \right).$$

For pseudo-real representations we can apply the constraint $\tilde{Q} = \pm\sigma_G Q$ to recover the Lagrangian

$$\mathcal{L}_{\text{hh}} = \int d^2\theta\, d^2\bar{\theta} \left(Q^\dagger e^V Q \right) \pm \sqrt{2}\ \text{Re} \int d^2\theta \left(Q^t \sigma_G^t\, \Phi\, Q \right)$$

for a single half-hypermultiplet. Here we rescaled $Q \to \frac{1}{\sqrt{2}} Q$ to give the kinetic term in the Lagrangian a canonical coefficient. We also used

$$\tilde{Q}^t e^{-V} \tilde{Q}^* = Q^t \sigma_G^t e^{-V} \sigma_G Q^* = Q^t e^{V^t} Q^* = Q^\dagger e^V Q\,,$$

since $\sigma_G^{-1} T \sigma_G = -T^t$ for $T \in \mathfrak{g}$. Since we found the Lagrangian $\mathcal{L}_{\text{hh}}$ by starting out with the Lagrangian $\mathcal{L}_{\text{fh}}$ for a full hypermultiplet and then applying the constraint $\tilde{Q} = \pm\sigma_G Q$, it is automatically invariant under $\mathcal{N} = 2$ supersymmetry.

Let us spell this out in some more detail. Substituting the constraint $\tilde{Q} = \pm\sigma_G Q$ in the $\mathcal{N} = 2$ supersymmetry equations yields two identical copies of the supersymmetry variations for the components of Q, which depend on all of the $\mathcal{N} = 2$ supersymmetry parameters. The Lagrangian $\mathcal{L}_{\text{hh}}$ is obviously invariant under these variations. The $SU(2)_R$ symmetry now acts on the vector of complex scalars $(q, \pm\sigma_G q^*)^t$.

As an example, the Lagrangian for the trifundamental $SU(2)$ half-hypermultiplet reads in components

$$\begin{aligned} \mathcal{L}_{\text{trif}} = &\int d^2\theta\, d^2\bar{\theta} \left(Q^*_{abc} e^{(V_1)^a{}_{a'}} Q^{a'bc} + Q^*_{abc} e^{(V_2)^b{}_{b'}} Q^{ab'c} + Q^*_{abc} e^{(V_3)^c{}_{c'}} Q^{abc'} \right) \\ &\pm \sqrt{2}\ \text{Re} \int d^2\theta \left(\epsilon^{bb'} \epsilon^{cc'} Q_{abc}\, \Phi^{aa'}\, Q_{a'b'c'} + \epsilon^{aa'} \epsilon^{cc'} Q_{abc}\, \Phi^{bb'}\, Q_{a'b'c'} + \epsilon^{aa'} \epsilon^{bb'} Q_{abc}\, \Phi^{cc'}\, Q_{a'b'c'} \right). \end{aligned} \tag{2.3}$$

We can obtain the bifundamental hyper by demoting one of gauge groups to a flavor group. From this perspective it is clear that the bifundamental has an enhanced $SU(2)$ flavor symmetry, as we already knew from general principles. We discuss these and other aspects of the $SU(2)$ trifundamental in detail in appendix A.

2.2 Instanton counting for half-hypermultiplets

We now turn to instanton counting for half-hypermultiplets. Also for this purpose it is convenient to use the description of a half-hypermultiplet as a constrained full hypermultiplet. Instanton counting for any $\mathcal{N} = 2$ gauge theory with full hypermultiplets is developed in [13, 8]

and spelled out in more detail in for example [14, 4]. It is performed by topologically twisting the $\mathcal{N} = 2$ gauge theory. The resulting instanton partition function is given by the integral

$$Z^{\text{inst}} = \sum_k q_{\text{UV}}^k \oint_{\mathcal{M}_k^G} e(\mathcal{V}) \tag{2.4}$$

over the ADHM moduli space of instantons $\mathcal{M}_k^G$ for the gauge group G and instanton number k, where the Euler class $e(\mathcal{V})$ encodes the matter content of the gauge theory. More precisely, the vector bundle $\mathcal{V}$ is equal to the space of solutions to the Dirac equation for the chosen matter representation in the self-dual instanton background.

Let us emphasize that the $SU(2)$ R-symmetry is essential for performing the topological twist. We identify the new Lorentz group of the twisted $\mathcal{N} = 2$ theory as

$$L' = SU(2)_L \times \text{diag}(SU(2)_R \times SU(2)_I) \ ,$$

where we denoted the R-symmetry group by $SU(2)_I$ to avoid confusion. After twisting the two complex scalars of a full hypermultiplet combine into a Weyl spinor

$$\Psi = (\psi_1, \psi_2) = (q, \tilde{q}^*) \ ,$$

i.e. the R-symmetry index turns into a spinor index. The matter part of the theory localizes to solutions of the Dirac equation [5]

$$(i\sigma^\mu \partial_\mu + \sigma^\mu A_\mu)\Psi = 0 \tag{2.5}$$

in the self-dual instanton background determined by the gauge field A, with $\sigma^\mu = (\mathbf{1}, i\sigma^i)$ (note that we are in Euclidean signature). These solutions form a vector bundle over the moduli space of self-dual instantons, localizing the path-integral to the integral over the moduli space of instantons (2.4). Actually computing the instanton partition function (2.4) can then be reduced to evaluating the equivariant index of the Dirac operator with respect to a torus action on the ADHM moduli space.

For half-hypermultiplets the twisting works similar, since we have established the R-symmetry invariance of the half-hypermultiplet Lagrangian. Let us start with a twisted full hypermultiplet. Since the R-symmetry indices of the scalars in the full hypermultiplet turn into spinor indices, we can again define the map

$$\Psi \mapsto \tau(\Psi) = \sigma_G \otimes \sigma_I \Psi^* \ . \tag{2.6}$$

[5] Although we write down an explicit form of the Dirac equation for a spinor transforming in the fundamental representation of a single gauge group, equation (2.5), as well as the following equations, should be read abstractly and can easily be adapted to hold in a generic setting.

As before, the matrices σ_G and σ_I act on the gauge and spinor indices, respectively. In particular,

$$\sigma_I \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} = \begin{pmatrix} -\psi_2 \\ \psi_1 \end{pmatrix}$$

The path integral of the half-hypermultiplet theory localizes onto solutions of the Dirac equation (2.5) that are invariant under τ.

The involution τ indeed maps solutions of the Dirac equation (2.5) to solutions, as can be seen from

$$(i\sigma^\mu \partial_\mu + \sigma^\mu A_\mu)\tau(\Psi) = -\sigma_G \otimes \sigma_I((i\sigma^\mu \partial_\mu + \sigma^\mu A_\mu)\Psi)^* = 0 \ , \tag{2.7}$$

where we have used $\sigma_I^{-1}\sigma^\mu \sigma_I = (\sigma^\mu)^*$ and $\sigma_G^{-1} A_\mu \sigma_G = -A_\mu^*$. We can thus find a basis of the space of solutions to the Dirac equation on which τ acts with eigenvalue ± 1. The relevant solutions for the single half-hypermultiplet are given by those basis elements which all have eigenvalue $+1$ (or all eigenvalue -1) under τ, and form a half-dimensional vector bundle over the moduli space $\mathcal{M}_G^k$ of self-dual instantons.



Figure 4: The solutions to the Dirac equation in a given representation of the gauge group form a vector bundle $\mathcal{V}$ over the ADHM moduli space $\mathcal{M}$. A pseudo-real representation induces a real structure τ on the vector bundle $\mathcal{V}$ that splits it into two copies $\mathcal{V} = \mathcal{V}_{\mathbb{R}} \oplus i\mathcal{V}_{\mathbb{R}}$. The relevant solutions for a half-hypermultiplet are either parametrized by $\mathcal{V}_{\mathbb{R}}$ or $i\mathcal{V}_{\mathbb{R}}$.

As an intermezzo, remember that the space of fermionic solutions to the Dirac equation in a pseudo-real representation always admits a real structure. It is not hard to see that the anti-linear involution τ in fact defines this real structure. So let us consider a basis of solutions on which τ acts with eigenvalues ± 1. Whereas for a theory with a hyper all solutions with eigenvalue $+1$ or -1 need to be taken into account, the theory with a half-hyper enforces a restriction to the solutions with eigenvalues either all $+1$ or all -1.

Let us name V the total vector space of solutions to the Dirac equation in a given instanton background. Then the real structure τ induces a splitting

$$V = V_R \oplus iV_R \ . \tag{2.8}$$

The vector space V_R (called the real form of τ) consists of solutions with eigenvalue $+1$, whereas iV_R consists of solutions with eigenvalue -1. Indeed, since τ is anti-linear, multiplying a solution $\Psi \in V_R$ by i yields a solution with eigenvalue -1. The real structure τ reduces the group of basis transformations acting on V from $U(d)$ to $SO(d)$, where d is the dimension of V. The action of $SO(d)$ leaves V_R invariant.

The two half-hypermultiplets that make up a hypermultiplet are defined by the two constraints $\tilde{Q} = \pm\sigma_G Q$. One half-hypermultiplet singles out the subspace $V_R \subset V$, and the other the subspace $iV_R \subset V$. So multiplying the solutions of the Dirac equation by i brings us from one half-hypermultiplet to the other.

Instanton partition functions for half-hypermultiplets

Let us summarize the above. Consider an $\mathcal{N} = 2$ gauge theory coupled to a full hypermultiplet in a pseudo-real representation of the gauge group. Its instanton partition function is given by equation (2.4). This is in fact equal to

$$Z^{\text{inst}}_{\text{fh}} = \sum_k q^k_{\text{UV}} \oint_{\mathcal{M}^G_k} e(\mathcal{V}_R \oplus i\mathcal{V}_R), \tag{2.9}$$

since the pseudo-real representation defines a real structure on the complex vector bundle $\mathcal{V}$ of solutions to the Dirac equation. The bundle $\mathcal{V}_R$ is an oriented real bundle, whose Euler class is defined as the Pfaffian (this is only non-trivial when the rank of the bundle is even). The Euler class of its complexification $\mathcal{V} = \mathcal{V}_R \oplus i\mathcal{V}_R$ can then be expressed as the square of the Euler class of $\mathcal{V}_R$,

$$e(\mathcal{V}_R \oplus i\mathcal{V}_R) = e(\mathcal{V}_R)^2. \tag{2.10}$$

This equality continues to hold for the equivariant Euler classes $e_{\mathbf{T}}(\mathcal{V})$ and $e_{\mathbf{T}}(\mathcal{V}_R)$, with respect to the torus action $\mathbf{T} = T_{a_k} \times T_{\phi_i} \times U(1)_{\epsilon_1,\epsilon_2}$ on the ADHM moduli space, where T_{a_k} is the torus of the gauge group, T_{ϕ_i} the torus of the dual group, and the action of $U(1)_{\epsilon_1,\epsilon_2}$ on $\mathbb{R}^4$ defines the Omega-background.

In other words, when the rank of $\mathcal{V}_R$ is even (*i.e.* when the complex Dirac index of the pseudo-real representation is even), the instanton partition function for a half-hyper theory localizes as

$$Z^{\text{inst}}_{\text{hh}} = \sum_k q^k_{\text{UV}} \oint_{\mathcal{M}^G_k} e(\mathcal{V}_R). \tag{2.11}$$

Since the involution τ commutes with the torus $\mathbf{T}$, we can compute the contribution of a half-hypermultiplet equivariantly by just taking the square-root of the product of weights for the full hypermultiplet theory.

In this manner we can compute the instanton partition function for the $Sp(1)$ trifundamental half-hypermultiplet and the $Sp(1) - SO(4)$ bifundamental half-hypermultiplet. [6] Notice that in both examples the Dirac index is even for any instanton number k. In section 5 we apply this scheme to evaluate instanton partition functions corresponding to $Sp(1)/SO(4)$ quiver gauge theories.

3. CFT building blocks for Sicilian quivers

Let us now discuss the building blocks that are needed for the AGT correspondence. In the correspondence for conformal $SU(2)$ quiver gauge theories hypermultiplets are given by punctures on the Gaiotto curve. Gluing the neighborhoods of two punctures to create a tube gauges the flavor symmetry group of the two hypermultiplets into an $SU(2)$ gauge group. The masses of the two hypermultiplets have to be opposite to perform the gluing, since they correspond to the residue of the Seiberg-Witten 1-form at the puncture. The masses then turn into the Coulomb parameters $\pm a$ of the $SU(2)$ gauge group after the gluing.

On the CFT side, hypermultiplets correspond to insertions of primary fields ϕ^i whose conformal weights are related to the masses of the hypermultiplets. A gauge group corresponds to inserting a complete set of descendants of a given primary field. We recall that an arbitrary Virasoro descendant ϕ_I at level N is given by a partition I of N by $\phi_I = \prod_j L_{-I_j}\phi$. For ease of notation we will also just write I for N. The projector on a particular representation that we insert can thus be written as

$$P_{\mathcal{H}_\phi} = \sum_{I,J} K^{-1}_{IJ} |\phi_J\rangle\langle\phi_I| \ , \tag{3.1}$$

where K^{-1} is the inverse of the Kac matrix $(K)_{IJ} = \langle\phi_I|\phi_J\rangle$. The modulus of the tube corresponds to the coupling of the gauge group. From this it is clear that if we decouple the gauge group by sending $q \to 0$ we recover the original expression for the ungauged theory, since the contributions of the descendants vanish and only the primary field survives.

The complete instanton partition function can thus be obtained from a pair of pants decomposition of the Gaiotto curve. Its building blocks are given by three-point functions containing one or more descendant fields, and the total expression is obtained by summing over all descendant fields in the channels. This sum corresponds to the sum over the fixed points in the instanton counting. For linear and cyclic quivers, the only building blocks needed are hypermultiplets in the fundamental and hypermultiplets in the bifundamental. The corresponding CFT expressions are three-point functions with one or two descendant fields.

[6] This derivation justifies the method used in [4] to compute the contribution of the $Sp(1) - SO(4)$ bifundamental half-hypermultiplet.



Figure 5: Decomposition of the sphere with six punctures into three-punctured spheres and tubes, and the corresponding conformal blocks.

For Sicilian quivers such as in figure 5, however, we also need hypers in the fundamental of three different gauge groups. The corresponding CFT building block should then be described by the three-point function with three descendant fields inserted,

$$\langle V(\phi_{I_1}^1, z_1)V(\phi_{I_2}^2, z_2)V(\phi_{I_3}^3, z_3)\rangle \ . \tag{3.2}$$

Here we have used the notation $V(\phi, z)$ for the the vertex operator corresponding to the field ϕ inserted at z. The weights of the fields ϕ^i are related to the Coulomb branch parameters $a_{1,2,3}$ of the three $SU(2)$ gauge groups involved. Choosing the insertion points z_i is quite subtle and affects the outcome, as we will now discuss.

3.1 Three-point functions

Let us start with a reminder about three-point functions and some of their properties. For three primary fields the three-point function is fixed up to a constant C_{123}. The coordinate dependence itself is fixed covariance under by Möbius transformations, *i.e.* the global conformal symmetry.

Local conformal symmetry allows us to compute three-point functions of arbitrary descendants of those primary fields as well. In principle, this is straightforward: the only thing needed is the OPE of the stress energy tensor $T(z)$ with the primary fields with itself. We can then use

$$\langle V(L_{-n}\phi, z)\dots\rangle = \oint_z dw\,(w-z)^{-n+1}\langle T(w)\,V(\phi, z)\cdots\rangle \tag{3.3}$$

to reduce the three-point function to contour integrals of the correlator of three primary fields and several energy stress tensors. This correlator is a meromorphic function on a Riemann surface and thus determined by its poles. We can thus consecutively eliminate the $T(z)$ by summing their OPEs with the other $T(w)$ and the primary fields, until we are left with just the three-point function of the primary fields. We can then evaluate the contour integral.

Though conceptually simple, in practice this procedure is quite cumbersome. Since most of the time we are interested in very specific values of z_i only, it can be more efficient to phrase the computation in terms of operators on the Hilbert space of a Virasoro representation. The operator-state correspondence tells us that

$$\lim_{z\to 0}\phi(z)|0\rangle = |\phi\rangle\,. \tag{3.4}$$

The corresponding bra state is given by the operator at infinity. More precisely, it is obtained from the ket state using the Möbius transformation $z \mapsto 1/z$:

$$\lim_{z\to 0}\langle 0|V(z^{-2L_0}e^{-\frac{1}{z}L_1}\phi, 1/z) = \langle\phi|\,. \tag{3.5}$$

The three-point function with primary fields at 0,1,∞ can then be computed as [7]

$$\langle\phi^1|V(\phi^2,1)|\phi^3\rangle = \langle\phi^1|\phi^2_{h_3-h_1}|\phi^3\rangle = C_{123}\,. \tag{3.6}$$

We can compute such three-point functions with descendant bra and ket states by commuting through all Virasoro operators using

$$[L_n,\phi_m] = (n(h-1)-m)\phi_{m+n} \tag{3.7}$$

for primary fields ϕ^2. If ϕ^2 is a descendant, then we first need to express it in terms of Virasoro operators and modes of the primary field, which we do by using the following expression for the $-N_1$ mode of a $V(L_{-N_2}\phi,z)$ [15]

$$\begin{aligned} V_{-N_1}(L_{-N_2}\phi) = &\sum_{l\geq 0}\binom{N_2-2-l}{l}L_{-N_2-l}V_{-N_1+N_2+l}(\phi) \\ &+(-1)^{N_2}\sum_{l\geq 0}\binom{N_2-2-l}{l}V_{-N_1-l+1}(\phi)L_{l-1}\,. \end{aligned} \tag{3.8}$$

Note that even though the sums are infinite, they reduce to finite sums when acting on any particular state.

[7] Strictly speaking we can only do this for $h_3-h_1\in\mathbb{Z}$. From general arguments we know however that the coefficients of the conformal block are given by rational functions in h_i and c. The expressions we obtain thus continue to be valid for arbitrary values of h.

Consider a theory with several identical gauge groups. One would expect that the partition function should be symmetric under suitable permutations of the gauge group. On the CFT side this means that the three-point function should be symmetric under permutations of the insertion points.

For instance, if a theory contains a hypermultiplet in the fundamental of two $SU(2)$ groups, then the three-point function must be symmetric under exchanging the two. This is indeed the case, as follows from

$$\langle \phi^1_{I_1} | V(\phi^2, 1) | \phi^3_{I_3} \rangle = \langle \phi^3_{I_3} | V(\phi^2, 1) | \phi^1_{I_1} \rangle \ . \tag{3.9}$$

To see that (3.9) indeed holds we can use the Möbius transformation $z \mapsto 1/z$. A general field transforms under a Möbius transformation γ as

$$V(\phi, z) \mapsto V\left((\gamma(z)')^{L_0} e^{\frac{\gamma''}{2\gamma'} L_1} \phi, \gamma(z) \right) \ . \tag{3.10}$$

From this we see that as long as ϕ^2 is a primary field, it does not pick up any correction terms from this transformation.

On the other hand, if we consider the case of a hyper in the fundamental of three gauge groups, we need to insert three descendants, and $V(\phi^2, 1)$ will no longer transform in such a simple way. The usual vertex is then no longer symmetric under permutations, as the Möbius transformations that exchange punctures introduce corrections. This means that the standard CFT vertex must correspond to a regularization scheme of the gauge theory which treats the gauge groups differently.

More generally, if we use any Möbius transformation to change the insertion points of a three-point function with descendants, then due to (3.10) we will pick up corrections. This means that the detailed expression for the three-point function greatly depends on the choice of insertion points z_i in (3.2). It turns out that these issues are less severe for asymptotically free theories. Let us therefore turn to those cases.

3.2 Partition function for the trifundamental coupling

Conformal blocks a priori correspond to conformal gauge theories, as the flavor symmetries always work out in such a way that there are four fundamental hypers per $SU(2)$ gauge group. We can however obtain asymptotically free theories by sending the mass of hypers to infinity and so decoupling them. More precisely, to decouple a hyper of mass m in the fundamental of a gauge group of coupling q we take

$$q \to \Lambda/m \ , \qquad m \to \infty \ . \tag{3.11}$$

Here Λ is the scale of the newly asymptotically free theory. In this way we can obtain any asymptotically free partition function from a conformal block.

Let us use the procedure outlined above to compute the partition function of a half-hypermultiplet in the trifundamental of $SU(2)$. We start out with conformal theory which corresponds to a sphere with six punctures (see figure 5), but decompose it in a symmetric (*i.e.* non-linear) way:

$$Z = \sum_{I_1,I_2,I_3} \sum_{J_1,J_2,J_3} \langle \phi^{m_1} | V(\phi^{m_2},1) | \phi^{a_1}_{I_1} \rangle \langle \phi^{m_3} | V(\phi^{m_4},1) | \phi^{a_2}_{I_2} \rangle \langle \phi^{m_5} | V(\phi^{m_6},1) | \phi^{a_3}_{I_3} \rangle \\ \times K^{-1}_{I_1 J_1} K^{-1}_{I_2 J_2} K^{-1}_{I_3 J_3} \langle \phi^{a_1}_{J_1} | V(\phi^{a_2}_{J_2},1) | \phi^{a_3}_{J_3} \rangle \, q_1^{I_1} q_2^{I_2} q_3^{I_3} \,. \quad (3.12)$$

Note that we have chosen more or less by fiat that the trifundamental vertex, *i.e.* the three-punctured sphere in the center of the decomposition, is given by the sphere with punctures at $0, 1, \infty$. In view of the remarks in the previous section the result is certainly not symmetric under permutation of the gauge groups. To obtain the asymptotically free theory, that is the result for a single half-hyper in the trifundamental, we apply (3.11). It turns out that the resulting expression *is* symmetric under permutations up to spurious terms (which we explain in a moment). It is moreover independent on the choice of punctures of the three-punctured sphere in the center of the decomposition, up to a simple rescaling of the couplings q.

This rather surprising result can be better understood when computing asymptotically free theories using Gaiotto states [11]. Such a state $|h, \Lambda\rangle$ is an eigenstate of the Virasoro mode L_1 with eigenvalue Λ,

$$L_1 |h,\Lambda\rangle = \Lambda |h,\Lambda\rangle \qquad L_n |h,\Lambda\rangle = 0 \quad n \geq 2 \,. \quad (3.13)$$

More concretely such as state can be written as a power series in Λ

$$|h,\Lambda\rangle = \sum_{n=0}^{\infty} \Lambda^n |v_n\rangle \,, \quad (3.14)$$

where $|v_0\rangle = |h\rangle$ and $|v_n\rangle$ is a specific linear combination of Virasoro descendants of $|h\rangle$ at level n. These states can then be used to compute instanton partition functions for asymptotically free $SU(2)$ theories. The norm of such a state for instance gives the instanton partition function of pure $SU(2)$ gauge theory. Both states in this norm originate from decoupling a pair of hypermultiplets in the conformal $SU(2)$ gauge theory. The conditions (3.13) come from the poles of the quadratic differential $\phi_2(z)$ on the Gaiotto curve. (See [16] for a proof that this is equivalent to the infinite mass limit.)

It is natural to use the same strategy also for multiple gauge groups. The $SU(2)$ trifundamental can be obtained by decoupling three pairs of hypers in the conformal $SU(2)$ gauge theory corresponding to the six-punctured sphere. We thus compute the three-point function

$$Z_{\text{CFT}} = \langle h_1, \Lambda_1 | V(|h_2, \Lambda_2\rangle, 1) | h_3, \Lambda_3 \rangle \,. \quad (3.15)$$

This gives indeed the same expression as the one we obtained above. Now we can also explain why (3.15) is invariant under permutation of the three gauge groups (up to some trivial factors). As usual we use a Möbius transformation γ to exchange the three insertion points. From (3.10) and (3.13) it follows that the Gaiotto state $|h, \Lambda\rangle$ transforms to

$$e^{\frac{\gamma''}{2\gamma'}\Lambda}(\gamma')^h|h, \gamma'\Lambda\rangle \ , \tag{3.16}$$

so that after a redefinition of Λ the two three-point functions only differ by a spurious prefactor. Since this holds for any Möbius transformation, the result is essentially independent of the insertion points.

We propose that (3.15) is equal to the instanton partition function of a half-hyper in the trifundamental representation of $SU(2)$ (up to a spurious factor [8]). Even though we did not compute this partition function directly, we can perform several consistency checks on (3.15). First note that it has indeed a proper $\mathcal{F}_g$ expansion, *i.e.* that it can be written

$$Z = \exp\left(\sum_{g\geq 0}\hbar^{2g-2}\mathcal{F}_g\right), \tag{3.17}$$

with no higher negative powers of $\hbar$ appearing. Second, (3.15) reduces correctly to the $SU(2)$ bifundamental when we decouple one of the gauge groups. Finally, when setting $\Lambda_2 = \Lambda_3$ it agrees with the partition function of a $Sp(1) - SO(4)$ gauge theory with a single hyper in the bifundamental (details of this check can be found in section 5).

4. Towards a 4d/2d correspondence for Sicilian quivers

The simplest way to define a conformal $\mathcal{N} = 2$ Sicilian $SU(2)$ quiver gauge theory is through its M-theory construction. Wrap two M5 branes on a Riemann surface with punctures $\mathcal{C}$. The quiver theory corresponding to a particular duality frame is obtained from a decomposition of $\mathcal{C}$ into pairs of pants. The punctures of $\mathcal{C}$ correspond to hypermultiplets, and the tubes connecting the different pants correspond to $SU(2)$ gauge groups whose microscopic coupling constants are given by the complex structure moduli of the tubes.

The building blocks are thus spheres with three punctures or tubes. There are three different configurations. The sphere with two punctures and one tube corresponds to two hypermultiplets in the fundamental. The sphere with one puncture and two tubes corresponds to a hyper in the bifundamental of the two $SU(2)$. Finally, as a new element, there is the the sphere with three tubes. It corresponds to a half-hyper in the trifundamental. Since the half-hyper is massless, it is natural not to have a puncture for it in this building block. See figure 6 for an example.

[8] In the following we define a spurious factor as a factor that does not depend on the Coulomb branch parameters and only contributes to the first terms of the genus expansion of the free energy.



Figure 6: Illustration of the correspondence between instanton partition functions of Sicilian $SU(2)$ quiver gauge theories and Virasoro conformal blocks on the corresponding Gaiotto curve for the six-punctured sphere. Each $SU(2)$ gauge group in the quiver is mapped to a tube in the Gaiotto curve, whereas $SU(2)$ matter is represented by three-punctured spheres.

Quivers with asymptotically free gauge groups can always be obtained from conformal theories by sending the mass of one of the hypers to infinity.

We can then compute the conformal block for this quiver in the following way. First, at every puncture insert a primary field whose conformal weight is given by the mass of the hyper in the usual way. Second, for every tube insert a projector

$$P_{\mathcal{H}_\phi} = \sum_{I,J} q^{|I|} K^{-1}_{IJ} |\phi_J\rangle\langle\phi_I|$$

onto the channel that corresponds to the Coulomb branch parameter of the $SU(2)$ gauge group. The bra and ket state of that projector are inserted in the respective building blocks. The problem thus reduces to computing various three point functions

$$\langle V(\phi^1_{I_1}, z_1) V(\phi^2_{I_2}, z_2) V(\phi^3_{I_3}, z_3)\rangle \ ,$$

of primary or descendant fields. As pointed out above, the subtlety lies in the choice of insertion the points z_i. For linear and cyclic quivers, all the building blocks have only one or two descendant fields inserted. Using the usual coordinates on the sphere or torus, the descendant fields are always inserted at 0 or ∞, that is as bra and ket states, and there is always a primary field inserted at 1. Using this prescription the conformal block agrees with the $SU(2)$ instanton partition function.

For trifundamental hypers the situation is more subtle. We can insert three descendant fields at the points $0, 1, \infty$, but in general the result will not agree with the instanton computation, since we are using a different parametrization of the moduli space. Once expressed in

IR variables, the results will agree. To put it another way, there will be a map between the moduli space coordinates and the microscopic gauge coupling that will make them agree. Or more geometrically, the CFT correlators define a unique object on the Seiberg-Witten curve, that is independent on the chosen parametrization of the complex structure moduli space of the Gaiotto curve.

The situation is much simpler for asymptotically free gauge groups. In this case the conformal block will agree with the instanton partition function immediately, and will be essentially independent of the choice of insertion points.

Comparison with Nekrasov partition function: one-loop factor

As we consider theories with $\mathcal{N} = 2$ supersymmetry, the full Nekrasov partition function has tree-level, one-loop and instanton contributions,

$$Z_{\text{Nek}} = Z_{\text{clas}}(\tau_{\text{UV}})\, Z_{\text{1-loop}}\, Z_{\text{inst}}(\tau_{\text{UV}})\,. \tag{4.1}$$

The 4d/2d correspondence relates the purely representation-dependent piece of the Liouville correlator on the Gaiotto curve, that is the the conformal block, to the instanton partition function of the corresponding gauge theory in the Omega-background. Adding the classical contributions to the instanton partition function is crucial for finding good properties under coordinate changes on the complex structure moduli space of the Gaiotto curve (we spell this out explicitly in section 5). The one-loop factor can be identified with the three point function of Liouville theory. More properly, the full conformal block on the Gaiotto curve should be identified with the Nekrasov partition function on S^4 [1].

Let us check that this agreement continues to hold for Sicilian quivers. The one-loop factor can be found as a four-dimensional boson-fermion determinant in the Omega-background. Equivalently, it may be obtained from the equivariant index of the Dirac operator in the instanton background (see appendix B). The resulting contribution for the (full) $SU(2)$ trifundamental hypermultiplet is

$$\begin{aligned} Z^{\text{2trif}}_{\text{1-loop}} &= \prod_{n,m=1}^{\infty} \prod_{i,j,k=1}^{2} \left(a_i + b_j + c_k + \frac{Q}{2} + n\epsilon_1 + m\epsilon_2 \right)^{-1} \\ &\propto \prod_{i,j,k=1}^{2} \Gamma_2 \left(a_i + b_j + c_k + \frac{Q}{2} \Big| \epsilon_1, \epsilon_2 \right), \end{aligned} \tag{4.2}$$

where we take the Coulomb branch parameters $a_i = \pm a$, $b_j = \pm b$ and $c_k = \pm c$ of the three $SU(2)$ gauge groups and $Q = \epsilon_1 + \epsilon_2$. The Barnes' double gamma function $\Gamma_2(x|\epsilon_1, \epsilon_2)$ regularizes the infinite product. The one-loop partition function for the $SU(2)$ trifundamental half-hypermultiplet is given by a square-root of the above expression.

Agreement with the three-point function of Liouville theory follows by the same argument as for linear quivers [1]. Namely, the numerator of the DOZZ formula for the Liouville three-point function contains the product

$$\prod_{i,j,k=1}^{2} \Gamma_2 \left(a_i + b_j + c_k + Q/2\right), \tag{4.3}$$

which equals the double trifundamental contribution in equation (4.2). Remember that the product (4.3) corresponds to the one-loop contribution of the Nekrasov partition function on S^4, which splits into a chiral and anti-chiral contribution on $\mathbb{R}^4$. Indeed, it is equal to the absolute value squared of the one-loop contribution for the $SU(2)$ trifundamental half-hyper, which for example can be written as

$$Z^{\text{trif}}_{\text{1-loop}} = \Gamma_2(a+b+c+Q/2)\Gamma_2(a+b-c+Q/2)\Gamma_2(a-b+c+Q/2)\Gamma_2(-a+b+c+Q/2).$$

5. Examples

In this section we test our proposal for extending the 4d/2d AGT correspondence to Sicilian quivers in the two examples illustrated in figure 7 and figure 8.



Figure 7: From a gauge theory perspective the $Sp(1) - SO(4)$ bifundamental, which is illustrated on the left, is equivalent to the $SU(2)$ trifundamental, which is illustrated on the right, once we identify two of the $SU(2)$ gauge couplings.

The quiver on the left in figure 7 consists of a single $Sp(1)$ gauge group and a single $SO(4)$ gauge group coupled by a bifundamental $Sp(1) - SO(4)$ half-hypermultiplet. It is equivalent to an $SU(2)$ Sicilian quiver gauge theory consisting of three $SU(2)$ gauge groups coupled by an $SU(2)$ trifundamental half-hypermultiplet, illustrated on the right in figure 7. The gauge couplings of both quivers are asymptotically free, so that the instanton partition function should agree directly with the CFT block (3.15) without any subtleties involving a choice of coordinates. We check that this is indeed the case up to order 3.

The quiver in figure 8 is a conformal $Sp(1) - SO(4)$ gauge theory with two bifundamental $Sp(1) - SO(4)$ half-hypermultiplets, which is equivalent to a conformal $SU(2)$ Sicilian quiver gauge theory with three $SU(2)$ gauge groups coupled by two trifundamental $SU(2)$ half-hypermultiplets. Since the gauge theory is conformal, the results will depend on the choice of



Figure 8: From a gauge theory perspective the cyclic $Sp(1) - SO(4)$ quiver, which is illustrated on the left, is equivalent to the genus 2 $SU(2)$ quiver, which is illustrated on the right, once we identify two of the $SU(2)$ gauge couplings.

complex structure on the Gaiotto curve, and on the instanton counting scheme. Our proposal tells us which CFT configuration to choose to give direct agreement with the $Sp(1) - SO(4)$ instanton partition function, and we check that this indeed works up to order 3.

The conformal $SU(2)$ gauge theory can alternatively be described in terms of a massless full $SU(2)$ trifundamental hyper. So we can find its instanton partition function as well using the more conventional $U(2)$ instanton counting scheme. [9] We check that if we use the $U(2)$ trifundamental instanton counting scheme or choose different coordinates in the conformal block the results do agree in the IR. This confirms the general philosophy outlined above.

On a more technical level, the instanton counting formulae for the $Sp(1) - SO(4)$ quiver gauge theories can be found in appendix B. They are given by a multiple contour integral of a meromorphic integrand. This integrand consists of building blocks, each piece coming from a component of the quiver gauge theory. The contributions for the gauge theory nodes were already found in [8, 14]. We find the contribution for the $Sp(1) - SO(4)$ bifundamental half-hypermultiplet as outlined in section 2. We also make a proposal the integrand for the full $SU(2)$ trifundamental hypermultiplet. To actually evaluate these contour integrands, that is to find which of the poles contribute and to compute their residues, is an elaborate process, which we will describe later on.

5.1 The $SU(2)$ trifundamental as a $Sp(1) - SO(4)$ bifundamental

In this section we compute the instanton partition function of the $Sp(1) - SO(4)$ quiver gauge theory with a single bifundamental half-hypermultiplet. The quiver diagram is given in figure 7 and the corresponding Gaiotto curve is illustrated in figure 9.

[9] Notice that when we turn on the mass of this hypermultiplet, the theory does not have a string embedding anymore. This implies that we cannot find a Gaiotto curve. The Seiberg-Witten curve does exist, nevertheless, and can for example be found through a semi-classical approximation of the instanton partition function. See [17] for a related discussion.



Figure 9: The Gaiotto curve for the $SU(2)$ trifundamental is a double covering of the Gaiotto curve for the $Sp(1) - SO(4)$ bifundamental. (The corresponding quiver diagrams are illustrated in figure 7.)

Computing the instanton partition function

The instanton partition function of this theory is given by

$$Z_{\text{inst}}(q_1, q_2) = \sum_{k_1,k_2} q_1^{k_1} q_2^{k_2}\, Z_{k_1,k_2} \tag{5.1}$$

with

$$Z_{k_1,k_2} = \oint \prod_{i=1}^{n_1} d\phi_i \prod_{j=1}^{k_2} d\psi_j\, z_{\text{vec},k_1}^{Sp(1)}(\phi)\, z_{\text{vec},k_2}^{SO(4)}(\psi)\, z_{\text{bifund},k_1,k_2}^{Sp(1)-SO(4)}(\phi,\psi). \tag{5.2}$$

Here q_1 and q_2 correspond to the exponentiated gauge couplings of the $Sp(1)$ and $SO(4)$ gauge group, respectively, and $k_1 = 2n_1 + \chi_1$. As mentioned above, the main problem is to find the correct prescription for the contour integral, and to evaluate the residues of the poles in question. In the case of ordinary $SU(N)$ quiver gauge theories, the poles of the integrand only come from the vector multiplet contribution, and can be labeled by colored Young diagrams. In the case at hand, however, the $Sp(1) - SO(4)$ bifundamental does introduce additional poles, so that evaluating the contour integral becomes much more complicated. More precisely, besides the poles coming from the $Sp(1)$ and the $SO(4)$ vector multiplet, there are also poles

$$\psi_j = \pm\epsilon_+ \quad (\text{when } k_2 \text{ is odd}) \tag{5.3}$$

$$\phi_i = \pm\psi_j \pm \epsilon_+ \tag{5.4}$$

from the $Sp(1) - SO(4)$ bifundamental. Note that these poles intertwine $Sp(1)$ poles and $SO(4)$ poles.

A priori the integrals in (5.2) are over the real axis. We need to make a choice in moving the poles away from the real axis and the closing the contour. The usual prescription is to move $\epsilon_{1,2} \mapsto \epsilon_{1,2} + i0$ and then close the contour in the upper-half plane. We use this convention to deal with the vector multiplet poles. For the bifundamental poles, however, we need to choose the opposite prescription $\epsilon_+ \mapsto \epsilon_+ - i0$. This recipe originates from the description of the poles for the massive full bifundamental hypermultiplet. Similarly to the pole prescription in the $\mathcal{N} = 4$ ADHM construction, we introduce two additional equivariant parameters $\epsilon_3 = -\mu - \epsilon_+$ and $\epsilon_4 = \mu - \epsilon_+$, which we assume to have positive imaginary parts. [10] To find the pole prescription for the bifundamental half-hypermultiplet, we just set the mass μ to zero (which identifies $\epsilon_3 = \epsilon_4$). We furthermore encounter poles of the form $n\epsilon_1 - m\epsilon_3$ with $n \in \frac{1}{2}\mathbb{N}$, $m \in \mathbb{N}$, which we also need to include. Our prescription is to take $\mathrm{Im}(\epsilon_\alpha - \epsilon_\beta) \gg 0$ if $\alpha > \beta$ as in the reference [19].

With this recipe we are set to evaluate the integral as the sum of pole residues. For each integration variable ϕ_i or ψ_j we have a precise prescription, so that we can proceed integral by integral. In practice it is useful to replace $\epsilon_+ \mapsto -\epsilon_3$ in the equations (5.3) and (5.4) to avoid any source of confusion. After identifying the additional poles coming from the bifundamental, we substitute back $\epsilon_3 \mapsto -\epsilon_+$ to evaluate the integral. Note that the unrefined partition function can only be obtained by setting $\epsilon_1 + \epsilon_2 = 0$ after we have performed the integration.

For the quiver gauge theory with a single $Sp(1) - SO(4)$ bifundamental half-hyper (see figure 7) the additional bifundamental poles start to contribute at instanton number $k = (k_1, k_2) = (2, 1)$. There are 12 new poles at this order. To get agreement with the conformal block (3.15) it is essential to include these extra poles. Interestingly, in the unrefined limit their contribution happens to vanish, so that instanton counting becomes much simpler.

Comparison with the three-point function (3.15)

We identify the parameters of the conformal field theory and gauge theory to be

$$\begin{aligned}
\Lambda_1 &= -\frac{q_1}{\epsilon_1\epsilon_2},\ \Lambda_2 = -\frac{q_2}{16\epsilon_1\epsilon_2},\ \Lambda_3 = \frac{q_2}{16\epsilon_1\epsilon_2} \\
h_1 &= \frac{1}{\epsilon_1\epsilon_2}\left(\frac{Q^2}{4} - a^2\right), \\
h_2 &= \frac{1}{\epsilon_1\epsilon_2}\left(\frac{Q^2}{4} - \left(\frac{b_1 + b_2}{2}\right)^2\right), \\
h_3 &= \frac{1}{\epsilon_1\epsilon_2}\left(\frac{Q^2}{4} - \left(\frac{b_1 - b_2}{2}\right)^2\right), \\
c &= 1 + \frac{6(\epsilon_1 + \epsilon_2)^2}{\epsilon_1\epsilon_2},
\end{aligned} \tag{5.5}$$

[10] See for example [18] for a detailed discussion of the $\mathcal{N} = 4$ ADHM construction.

where c is the central charge and h_i are the conformal weights of the vertex operators. Comparing this with the three-point function (3.15) we find

$$\boxed{Z_{\text{inst}}(q_1, q_2) = Z_{\text{CFT}}(q_1, q_2) Z^{(1)}_{\text{spur}}\,,} \tag{5.6}$$

up to $k = (k_1, k_2) = (2, 2)$, with the spurious factor

$$Z^{(1)}_{\text{spur}} = \exp\left(\frac{q_1}{2\epsilon_1\epsilon_2}\right) \exp\left(\frac{q_2}{8\epsilon_1\epsilon_2}\right)\,.$$

5.2 Genus two quiver through $Sp(1) - SO(4)$ instanton counting

The Sicilian quiver theory for genus two Gaiotto curve is an $SU(2)^3$ theory with two trifundamental half-hypers. We can also view it as a $Sp(1) - SO(4)$ theory with two bifundamental half-hypers (see figure 8). As illustrated in figure 10, the corresponding Gaiotto curve is a torus with two punctures. The genus two curve is the double cover of the genus one curve with two branch points.



Figure 10: The Gaiotto curve for the $SU(2)$ genus 2 quiver is a double covering of the Gaiotto curve for the $Sp(1) - SO(4)$ cyclic quiver. (The quiver diagrams are illustrated in figure 8.)

The partition functions will depend on the choice of coordinates of the Riemann surface, because the gauge theory is conformal. Using our proposal for the $Sp(1) - SO(4)$ AGT correspondence, we can find a choice of coordinates on the complex moduli space of the genus two Gaiotto curve that matches the instanton partition function with the conformal block directly, without a UV-UV map.

Computing the instanton partition function

Since a single full hypermultiplet can be obtained by combining two half-hypermultiplets, there are two different ways to compute the instanton partition function: either as a single full bifundamental hyper or as two bifundamental half-hypers. Using the first method we start from the massive full hyper. The term at order (k_1, k_2) is given by

$$Z^{(1)}_{k_1,k_2} = \oint \prod_{i=1}^{n_1} d\phi_i \prod_{j=1}^{k_2} d\psi_j \, z^{Sp(1)}_{\text{vec},k_1}(\phi, a) \, z^{SO(4)}_{\text{vec},k_2}(\psi, b) \, z^{Sp(1),SO(4)}_{\text{2bif},k_1,k_2}(\phi, \psi, \mu). \tag{5.7}$$

where the explicit form of the integrand is given in appendix B.

We can evaluate the contour integral using the two equivariant parameters, $\epsilon_3 = -\mu - \epsilon_+$ and $\epsilon_4 = \mu - \epsilon_+$. The massive full bifundamental introduces the additional poles

$$\psi_j = \pm\epsilon_3, \pm\epsilon_4 \quad (\text{when } k_2 \text{ is odd}) \tag{5.8}$$

$$\phi_i = \pm\psi_j \pm \epsilon_3 \tag{5.9}$$

$$\phi_i = \pm\psi_j \pm \epsilon_4. \tag{5.10}$$

Compared to the single bifundamental half-hyper in the previous example the additional parameter ϵ_4 introduces extra poles. For example, there are in total 28 new poles with non-vanishing residues at $k = (1, 2)$. Again the contribution from these new poles happens to vanish in the unrefined limit, whereas it is crucial to include them in the refined setup.

Using the second method we start with the contour integral corresponding to two massless bifundamental half-hypers

$$Z^{(2)}_{k_1,k_2} = \oint \prod_{i=1}^{n_1} d\phi_i \prod_{j=1}^{k_2} d\psi_j \, z^{Sp(1)}_{\text{vec},k_1}(\phi, a) \, z^{SO(4)}_{\text{vec},k_2}(\psi, b) \left(z^{Sp(1),SO(4)}_{\text{bif},k_1,k_2}(\phi, \psi) \right)^2, \tag{5.11}$$

where the explicit form of each of the building blocks is given in appendix B. The poles of this integral are simpler to enumerate, since there is just one new equivariant parameter $\epsilon_3 = -\epsilon_+$. There are 10 new poles with non-vanishing residues at $k = (1, 2)$. As they should, both computations indeed give the same result once we set the mass $\mu = 0$.

CFT computation

Let us now compute the conformal block for the genus two surface. The most straightforward guess is to imitate (3.12) by taking

$$Z = \sum_{I_1,I_2,I_3} \sum_{J_1,J_2,J_3} K^{-1}_{I_1J_1} K^{-1}_{I_2J_2} K^{-1}_{I_3J_3} \langle \phi^{a_1}_{I_1} | V(\phi^{a_2}_{I_2}, 1) | \phi^{a_3}_{I_3} \rangle \langle \phi^{a_1}_{J_1} | V(\phi^{a_2}_{J_2}, 1) | \phi^{a_3}_{J_3} \rangle \, q_1^{I_1} q_2^{I_2} q_3^{I_3}. \tag{5.12}$$

By the general remarks above, this expression corresponds to a particular parametrization of the moduli space of the genus two surface. Presumably there should be a corresponding

regularization scheme on the gauge theory side. In particular, the conformal block (5.12) should agree with any instanton computation in the IR.

We want to do a bit better than that however: we want to find an expression which agrees with our instanton computation of the genus two surface as a cyclic $Sp(1) - SO(4)$ quiver in the UV. The AGT correspondence for Sp/SO quivers was worked out in [4]. Let us briefly summarize the relevant facts.

The Gaiotto curve of the cyclic Sp/SO theory is a torus with a $\mathbb{Z}_2$ branch cut running between two branch points. The double cover of this curve is the genus two curve where two of the moduli are equal, see figure 10. The $\mathcal{W}$-algebra of the theory is a double copy of the Virasoro algebra, where the $\mathbb{Z}_2$-twist exchanges the two copies. The conformal block of this configuration on the torus with total modulus $q_1^2 q_2$ is given by

$$\mathrm{Tr}\left[\sigma(1) P_{a_1} \sigma(q_1^2) P_{a_2,a_3} (q_1^2 q_2)^{L_0}\right] . \tag{5.13}$$

Here σ is the $\mathbb{Z}_2$-twist vacuum, and we take the branch cut to go from $\sigma(1)$ through P_{a_1} to $\sigma(q_1^2)$. P_{a_1} is the projector onto the twisted representation coming from the primary field ϕ^1. As the primary field ϕ^1 transforms in a twisted representation, it is indeed characterized by a single parameter a_1. On the other hand, P_{a_2,a_3} is the projector onto the untwisted representation characterized by two parameters a_2 and a_3.



Figure 11: The map $\gamma(z)$ relates the $\mathcal{W}$-block on the twice punctured torus (which computes the double $Sp(1) - SO(4)$ instanton partition function) to a Virasoro block on its double cover, a genus two curve. More precisely, we first cut open the torus along the $SO(4)$ tube and insert a complete basis of states $V^{a_2,a_3}_{I_2,I_3}$ in the untwisted representation labeled by a_2 and a_3. Then we map this onto a genus two surface using the map γ, and insert a complete basis of states $V^{a_1}_{I_1}$ in the Virasoro representation labeled by a_1.

$$\langle V^{a_2,a_3}_{I_2,I_3}(\infty)\sigma(1)P_{a_1}\sigma(q_1^2)V^{a_2,a_3}_{J_2,J_3}(0)\rangle$$

$$\downarrow z \mapsto \gamma(z)$$

$$\langle \tilde{V}^{a_2}_{I_2}(1)\tilde{V}^{a_3}_{I_3}(-1)|V^{a_1}_{I_1}\rangle\langle V^{a_1}_{J_1}|\tilde{V}^{a_2}_{J_2}(q_1)\tilde{V}^{a_3}_{J_3}(-q_1)\rangle$$

$$\swarrow \qquad\qquad \searrow z \mapsto q_1/z$$

$$\langle \tilde{V}^{a_2}_{I_2}(1)\tilde{V}^{a_3}_{I_3}(-1)|V^{a_1}_{I_1}\rangle \qquad\qquad \langle \tilde{V}^{a_2}_{J_2}(1)\tilde{V}^{a_3}_{J_3}(-1)|V^{a_1}_{J_1}\rangle$$

Table 1: Sequence of maps used for computing the cyclic $Sp(1) - SO(4)$ quiver.

.

Computing correlators of twisted representations can be done by going to the cover of the surface. Here, the $\mathcal{W}$-algebra on the cover is a single copy of the Virasoro algebra, and the problem reduces to the computation of the standard conformal block on the genus two surface. Although conceptually straightforward, this procedure leads to some technical subtleties. The main problem is that we apply the cover map to compute correlators with Virasoro descendants, which leads to correction terms. Let us therefore spell out precisely the cover map and the ensuing correlation functions.

To map the four point function (5.13) to the cover, we use the map γ

$$z \mapsto \gamma(z) = \pm\sqrt{\frac{z-q_1^2}{z-1}}\ , \tag{5.14}$$

where the sign determines the branch of the cover. This map indeed has branch points at $z = 1$ and $z = q_1^2$, and it maps the operators at 0 and ∞ to $\pm q_1$ and ± 1. This is illustrated in figure 11. In a second step, we want to reduce everything to standard building blocks, that is three-point functions on the sphere with operators inserted at $1, -1$ and 0.

The total sequence of maps is shown in table 1. [11] Introducing the notation

$$\mathbf{C}^{a_2,a_3;a_1}_{I_2,I_3,I_1}(q_1) = \langle \tilde{V}^{a_2}_{J_2}(1)\tilde{V}^{a_3}_{J_3}(-1)|V^{a_1}_{J_1}\rangle\, q_1^{2(h_{a_2}+h_{a_3}+I_2+I_3)},$$

the conformal block is given by

$$Z_{\text{CFT}} = \sum_{I_i,J_i} K^{-1}_{I_1J_1}K^{-1}_{I_2J_2}K^{-1}_{I_3J_3}\mathbf{C}^{a_2,a_3;a_1}_{I_2,I_3,I_1}(q_1)\mathbf{C}^{a_2,a_3;a_1}_{J_2,J_3,J_1}(q_1)q_1^{I_1}q_2^{I_2+I_3}. \tag{5.15}$$

[11] The $\tilde{V}(\pm q_1)$'s in the second line of table 1 are slightly different from the $\tilde{V}(\pm 1)$'s, which are defined in equation (5.17).

Note that the three-point functions $\mathbf{C}(q_1)$ depends on q_1 in a non-trivial way. They are obtained by acting with q_1-dependent coordinate transformations, so that the vertex operators $\tilde{V}$ depend on q_1. Since (5.14) is not a Möbius transformation, we need the following generalization [20] of (3.10):

$$V(\phi, z) \mapsto V\left(\left[\prod_{n=1}^{\infty} \exp\left(\frac{T_n(z)}{f'(z)^n} L_n\right)\right] f'(z)^{L_0}\phi, f(z)\right), \tag{5.16}$$

which holds for z away from singular points. Here we take all products to go from left to right. The functions $T_n(z)$ are defined recursively. The first two are given by

$$T_1(z) = \frac{f''(z)}{2f'(z)}, \qquad T_2(z) = \frac{1}{3!}\left(\frac{f'''(z)}{f'(z)} - \frac{3}{2}\left(\frac{f''(z)}{f'(z)}\right)^2\right).$$

For the transformations in (5.15) the new vertex operators are given by

$$\tilde{V}_I^a(\pm 1) = \left(\pm\frac{1-q_1^2}{2q_1^2}\right)^{h_a+I} V\left(\exp\left[\pm\frac{3+q_1^2}{2(1-q_1^2)} L_1\right] \exp\left[\frac{1}{4}L_2\right] \cdots \phi_I^a, \pm 1\right), \tag{5.17}$$

where the dots signify exponential factors involving higher L_n.

When we compare the genus two conformal block (5.15) with the instanton counting result, we find that they indeed agree up to order $k = (1, 2)$:

$$\boxed{Z_{\text{inst}}(q_1, q_2) = Z_{\text{CFT}}(q_1, q_2) Z^{(2)}_{\text{spur}},} \tag{5.18}$$

where the spurious factor

$$Z^{(2)}_{\text{spur}} = 1 - \frac{3\left(\epsilon_1+\epsilon_2\right)^2}{8\left(\epsilon_1\epsilon_2\right)} q_2 + \frac{3\left(\epsilon_1+\epsilon_2\right)^2\left(3\epsilon_1^2 - 2\epsilon_1\epsilon_2 + 3\epsilon_2^2\right)}{128\epsilon_1^2\epsilon_2^2} q_2^2 + \ldots$$

does not depend on physical parameters.

5.3 Alternative prescriptions for the genus two quiver

In the previous example we chose the coordinates of the genus two Gaiotto curve in such a way that we obtained direct agreement between the conformal block and the $Sp(1) - SO(4)$ instanton counting. If we choose different coordinates, or if we use a different instanton counting scheme, then the result will be different. In the infrared, however, all versions should agree. Put differently, we should be able to find a map between the UV couplings that make two results agree. Let us show that this philosophy is correct in two examples.

Comparison using a different conformal block

First, we can use the "naive" conformal block (5.12) to compute the genus two correlator. [12] We indeed find agreement between the $Sp(1)-SO(4)$ instanton partition function and this genus two conformal block using the UV-UV map

$$\begin{aligned}
\tilde{q}_1 &= q_1 - \frac{1}{4}q_1q_2 - \frac{1}{2}q_1^2 - \frac{1}{16}q_1q_2^2 + \frac{1}{4}q_1^2q_2 + \frac{3}{16}q_1^3 + O(q^4),\\
\tilde{q}_2 &= \frac{1}{16}q_2 - \frac{1}{16}q_1q_2 + \frac{3}{128}q_2^2 - \frac{1}{128}q_1q_2^2 + \frac{3}{128}q_1^2q_2 + \frac{55}{4096}q_2^3 + O(q^4),\\
\tilde{q}_3 &= \frac{1}{16}q_2 + \frac{1}{16}q_1q_2 + \frac{3}{128}q_2^2 + \frac{1}{128}q_1q_2^2 + \frac{3}{128}q_1^2q_2 + \frac{55}{4096}q_2^3 + O(q^4).
\end{aligned} \tag{5.19}$$

To compare the instanton partition functions with different parametrization of couplings, it is important to include tree-level pieces.[13] Similarly we also need to include the tree-level piece of the conformal block. In the case at hand we have

$$Z^{Sp-SO}_{\text{tree}}(q) = q_1^{-\frac{a^2}{\epsilon_1\epsilon_2}}\left(\frac{q_2}{16}\right)^{-\frac{1}{2\epsilon_1\epsilon_2}(b_1^2+b_2^2)} \quad \text{and} \quad Z^{\text{CFT}}_{\text{tree}}(\tilde{q}) = \tilde{q}_1^{-\frac{a_1^2}{\epsilon_1\epsilon_2}}\,\tilde{q}_2^{-\frac{a_2^2}{\epsilon_1\epsilon_2}}\,\tilde{q}_3^{-\frac{a_3^2}{\epsilon_1\epsilon_2}}\,.$$

Note that these two factors are related by the identifications (5.5) and the mappings $a_2 \to \frac{1}{2}(b_1+b_2)$ and $a_3 \to \frac{1}{2}(b_1-b_2)$. Using the UV-UV mapping (5.19) and the above identification of Coulomb parameters, we find that the ratio of Z_{Nek} and Z_{CFT} is given by the spurious factor

$$Z^{(3)}_{\text{spur}} = 1 - \frac{(\epsilon_1+\epsilon_2)^2}{8\epsilon_1\epsilon_2}q_1 - \frac{(\epsilon_1+\epsilon_2)^2}{4\epsilon_1\epsilon_2}q_2 + \frac{(\epsilon_1+\epsilon_2)^2(\epsilon_1^2+3\epsilon_1\epsilon_2+\epsilon_2^2)}{32\epsilon_1^2\epsilon_2^2}q_1q_2 + \cdots\,.$$

Comparison using a different method of instanton counting

As another test of this philosophy on the gauge theory side, we can compute the instanton using the $U(2)$ instanton counting scheme. This is possible since two $SU(2)$ trifundamental half-hypers combine into a single massless $SU(2)$ full trifundamental hyper. Apart from a non-trivial UV-UV mapping, we expect the $U(2)$ instanton and the CFT computation to differ by a non-trivial $U(1)$ factor. This $U(1)$ factor should not depend on any of the Coulomb parameters, after we enforce the tracelessness condition on all three gauge groups.

Since the trifundamental hyper is in the fundamental representation of the three gauge groups $SU(2)_A \otimes SU(2)_B \otimes SU(2)_C$, we find the contour integrand by considering the tensor product

$$\mathcal{E}_A \otimes \mathcal{E}_B \otimes \mathcal{E}_C \otimes \mathcal{L}$$

[12] We additionally checked that genus two correlator is independent on the choice of internal punctures up to a UV-UV mapping.

[13] One-loop factors are not relevant since they do not involve the gauge couplings.

of the three $U(2)$ universal bundles $\mathcal{E}_{A/B/C}$ over the product of ADHM moduli spaces $\mathcal{M}_{U(2)_A} \times \mathcal{M}_{U(2)_B} \times \mathcal{M}_{U(2)_C} \times \mathbb{R}^4$ with the half-canonical bundle $\mathcal{L}$ over $\mathbb{R}^4$. Recall that the restriction of each universal bundle to a self-dual connection $\mathcal{A}$ is just the corresponding instanton bundle $\mathcal{E}|_{\mathcal{A}} = E$ over $\mathbb{R}^4$ (see for example [4] for more details). The equivariant weights contributing to the Euler class $e_{\mathbf{T}}(\mathcal{V})$ of the bundle $\mathcal{V}$ of Dirac zero modes can then be found from the equivariant Dirac index

$$\mathrm{Ind}_{\mathbf{T}}(\not{D})^{g=2}_{U(2)^3} = \int_{\mathbb{C}^2} \mathrm{Ch}_{\mathbf{T}}(\mathcal{E}_A \otimes \mathcal{E}_B \otimes \mathcal{E}_C \otimes \mathcal{L})\, \mathrm{Td}_{\mathbf{T}}(\mathbb{C}^2). \tag{5.20}$$

The resulting contour integrand can be found in appendix B. For up to two instantons, it reproduces the partition functions for the bifundamental. For three non-zero instanton numbers the evaluation of the contour integral becomes tricky, because, unlike for $U(2)$ bifundamentals, many additional poles appear. It would be interesting to find an elegant prescription for the additional poles that yields agreement with the CFT.

We can still compare the instanton partition function up to second order. We find that up to this order it agrees with the conformal block (5.12), when we use the map

$$\begin{aligned}
\tilde{q}_A &= q_A + 2q_A^2 q_B + 6q_A q_B q_C + \mathcal{O}(q^4) \\
\tilde{q}_B &= q_B + 2q_A q_B + 2q_B q_C + 8q_A q_B q_C + 3q_A^2 q_B - 2q_A q_B^2 + 2q_B^2 q_C + 3q_B q_C^2 + \mathcal{O}(q^4) \\
\tilde{q}_C &= q_C + 2q_B q_C + 6q_A q_B q_C + 3q_B^2 q_C + 2q_B q_C^2 + \mathcal{O}(q^4)
\end{aligned}$$

between gauge coupling constants $\tilde{q}_{A/B/C}$ and complex structure parameters $q_{A/B/C}$, and up to a (unrefined) $U(1)$ factor

$$Z_{\mathrm{U}(1)} = 1 + q_A q_B + q_B q_C + q_C q_A + \cdots \tag{5.21}$$

Again, we need to include the classical contributions here. The non-trivial UV-UV mapping is expected since the conformal three-point function only reduces directly to a bifundamental contribution when two of its punctures are set at the positions 0 and ∞, and the primary vertex operator is inserted at 1.

6. Discussion

Let us briefly summarize our results and discuss some open questions. In this paper we extended the AGT correspondence to Sicilian quivers. To do this we first pointed out that the instanton partition function corresponding to the $SU(2)$ trifundamental cannot be found using the more conventional $U(2)$ instanton counting scheme, but should be computed using either an $Sp(1)^3$ or an $Sp(1) - SO(4)$ instanton counting scheme. We also argued that since the former does not have a string theory embedding, it is more natural to use the latter to find a corresponding CFT configuration.

As one of our main result we found that the instanton partition function for the $SU(2)$ trifundamental interaction is equal to the three-point function of Gaiotto states

$$\boxed{\langle h_1, \Lambda_1 | V(|h_2, \Lambda_2\rangle, 1) | h_3, \Lambda_3 \rangle.} \tag{6.1}$$

We have verified that it satisfies various consistency checks, such as symmetry under exchange of the gauge groups and reduction to the $SU(2)$ bifundamental. Furthermore, it agrees with the instanton partition function for the $Sp(1) - SO(4)$ bifundamental.

We have also proposed a construction for the Virasoro conformal block corresponding to any asymptotically free or conformal generalized $SU(2)$ quiver gauge theory. This proposal again passes all consistency checks, such as having a a proper genus-expansion in the parameter $\hbar$, and it has the expected properties in terms of UV vs. IR parameters. These are non-trivial properties for the conformal blocks, and it would be interesting to find a CFT explanation. Secondly, we have checked the proposal against $Sp(1) - SO(4)$ instanton counting in several Sicilian examples. To get an exact agreement of the instanton partition functions with the conformal blocks, we have used the string embedding of the $Sp(1) - SO(4)$ gauge theories to find the correct parametrizations of the conformal blocks.

There remain several interesting questions. It would for example be insightful to find a (geometric, or gauge theoretic) explanation for the asymmetry of the three-point function with three descendants. Relatedly, one can try to come up with a prescription for the parametrization of the conformal blocks dual to conformal Sicilian quivers that agrees on the nose with the instanton partition function. One can also wonder whether it is possible to perform instanton counting for any choice of coordinates on the complex structure moduli space. Why does instanton counting choose the particular parametrization it chooses? It would furthermore be interesting to look for a CFT object on the Seiberg-Witten curve that agrees with the IR partition function.

From the instanton counting perspective it is also curious that every instanton partition function can be decomposed into interactions with three or fewer gauge groups. This translates to the statement that any cohomology class on the instanton moduli space can be written as a product of only a few elementary classes. It would be interesting to understand this better, and necessary for a complete understanding of instanton counting for Sicilian quiver theories.

An exciting extension would be a verification of our proposal through geometric engineering [21] and the (refined) topological vertex [22]. It is not even obvious that the toric diagrams corresponding to Sicilian quivers indeed have the correct description under the decoupling of gauge groups. Furthermore, in this setting gauge groups are engineered using combinations of D4 and NS5-branes. This is very unusual from the perspective of geometric engineering, and requires a better understanding. Another fruitful direction would be the inclusion of BPS operators in these Sicilian gauge theories. Finally, a full verification of our proposal requires a

more detailed study of instanton counting for the $Sp(1)^3$ trifundamental half-hypermultiplet [23]. [14]

Acknowledgments

We would like to thank Fernando Alday, Alexander Braverman, Andrew Dancer, Nikita Nekrasov, Sara Pasquetti, Vasily Pestun, Yuji Tachikawa for enlightening discussions and correspondence. LH would like to thank the support of Sergei Gukov and the hospitality of the Mathematics Institute of the University of Oxford. The work of LH is supported by a NWO Rubicon fellowship and by NSF grant PHY-0757647. The work of CAK is supported by a John A. McCone Postdoctoral Fellowship. This work is in addition supported in part by the DOE grant DE-FG03-92-ER40701.

Appendix

A. More about the trifundamental half-hypermultiplet

In this appendix we explicitly show the reduction of the Lagrangian for the $SU(2)$ trifundamental half-hypermultiplet to bifundamental and fundamental hypermultiplets when we Higgs one or more of the $SU(2)$ gauge groups. Before starting this argument, let us quickly remind ourselves about flavor symmetry enhancement for (pseudo-)real representations.

Flavor symmetry enhancement

Let us briefly explain a way to understand the enhancement of flavor symmetries for matter transforming in a (pseudo-)real representation. Although this is not of direct importance for this paper, it will be useful as background and in the following.

First of all, recall the familiar statement of flavor symmetry enhancement. The flavor symmetry group of N hypers in a real representation of the gauge group is enhanced from $U(1)^N$ to $Sp(N)$, whereas the flavor symmetry of N hypers in a pseudo-real representation is enhanced to $SO(2N)$. The cases that are important for us is the single hyper in the bifundamental of $SU(2)$ which has enhanced flavor symmetry $Sp(1) = SU(2)$, and two hypers in the fundamental which enhance to $SO(4) = SU(2) \times SU(2)$.

A pseudoreal representation is characterized by an antilinear map σ_G such that $\sigma_G^2 = -1$. This map corresponds to the complex conjugation, so that in case of the fundamental or

[14] Starting from the $Sp(1)$ instanton formalism of [8, 14] and using the tools developed in this paper, we find an expression for the contour integral which has odd properties. It is quite possible that these may be caused by the lack of a string theory embedding of the $Sp(1)^3$ trifundamental half-hypermultiplet.

adjoint representation $\sigma_G^{-1} T \sigma_G = T^* = -T^t$ for $T \in \mathfrak{g}$. For real representations the only difference is that $\sigma_G^2 = 1$. Note that σ_G is automatically unitary.

The basic idea behind the enhancement is that the N-dimensional 'flavor vector' Q_i is enlarged to a $2N$-dimensional vector $(Q_i, \sigma_G \tilde{Q}_i)$ (which still is in the representation R of the gauge group). What needs to be shown is that the terms in the Lagrangian are invariants of $SO(2N)$ or $Sp(N)$.

For a single hypermultiplet in the fundamental representation of $SU(2)$, which is pseudo-real, the kinetic term in the Lagrangian $\mathcal{L}_{\text{fh}}$ can be rewritten as

$$\begin{aligned} Q^\dagger e^V Q + \tilde{Q}^t e^{-V} \tilde{Q}^* &= Q^\dagger e^V Q + \tilde{Q}^\dagger e^{-V^t} \tilde{Q} \\ &= Q^\dagger e^V Q + \tilde{Q}^\dagger \sigma_G^{-1} e^V \sigma_G \tilde{Q} \\ &= \left(Q^\dagger,\, \tilde{Q}^\dagger \sigma_G^\dagger \right) e^V \begin{pmatrix} Q \\ \sigma_G \tilde{Q} \end{pmatrix} . \end{aligned} \tag{A.1}$$

The Yukawa coupling in the Lagrangian $\mathcal{L}_{\text{fh}}$ is proportional to

$$\begin{aligned} 2\, \tilde{Q}^t \Phi Q &= Q^t \Phi^t \tilde{Q} + \tilde{Q}^t \Phi Q \\ &= Q^t \sigma_G \Phi \sigma_G \tilde{Q} + \tilde{Q}^t \Phi Q \\ &= \left(Q^t,\, \tilde{Q}^t \sigma_G^t \right) \sigma_G \Phi \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} Q \\ \sigma_G \tilde{Q} \end{pmatrix} . \end{aligned} \tag{A.2}$$

In both cases we used the fact $\sigma_G^{-1} T \sigma_G = -T^t$ for $T \in \mathfrak{g}$.

To see that (A.2) is an $SO(2)$ invariant, we can make a change of basis to $Q^\pm = Q \pm i\sigma_G \tilde{Q}$. The enhanced flavor group $SO(2)$ then acts in the fundamental on this new basis, and both (A.1) and (A.2) are the standard diagonal invariants. This argument generalizes in a straightforward way to an arbitrary number of hypers Q_i.

A similar argument works for real representations, such as for the bifundamental in $SU(2)_A \times SU(2)_B$. In this case the kinetic term can written exactly in the form (A.1) as well. The Yukawa term picks up a minus sign, due to the fact that now $\sigma_G^2 = 1$, so that the invariant is found by replacing

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \mapsto \begin{pmatrix} 0 & \mathbf{1}_N \\ -\mathbf{1}_N & 0 \end{pmatrix}$$

in (A.2). The Lagrangian is thus indeed invariant under $Sp(N)$. (Note that by $Sp(N)$ we actually mean $USp(N) = U(2N) \cap Sp(2N, \mathbb{C})$ here.)

Reduction of the $SU(2)$ trifundamental half-hypermultiplet

We first show that the Lagrangian for the $SU(2)$ trifundamental half-hyper reduces to that of a massive $SU(2)$ bifundamental hyper when one of the three $SU(2)$ gauge groups is

Higgsed. Start with the Yukawa terms

$$W = \epsilon^{bb'}\epsilon^{cc'}Q_{abc}\Phi_1^{aa'}Q_{a'b'c'} + \epsilon^{aa'}\epsilon^{cc'}Q_{abc}\Phi_2^{bb'}Q_{a'b'c'} + \epsilon^{aa'}\epsilon^{bb'}Q_{abc}\Phi_3^{cc'}Q_{a'b'c'} \tag{A.3}$$

in the Lagrangian of the $SU(2)$ trifundamental half-hyper (as derived in section 2). The first gauge group is Higgsed by setting $(\Phi_1)^a{}_{a'} = m_1(\sigma_3)^a{}_{a'}$, so that the superpotential W reduces to

$$W = m_1(\sigma_3)^{a'}{}_a Q^{abc}Q_{a'bc} - \epsilon^{cc'}Q^a{}_{bc}\Phi_2^{bb'}Q_{ab'c'} - \epsilon^{bb'}Q^a{}_{bc}\Phi_3^{cc'}Q_{ab'c'}. \tag{A.4}$$

When identifying

$$\begin{aligned} Q_{bc} &\equiv Q_{1bc} = -Q^2{}_{bc} \\ \tilde{Q}_{bc} &\equiv Q_{2bc} = Q^1{}_{bc} \end{aligned} \tag{A.5}$$

in equation (A.4), we indeed recover the Yukawa terms

$$W = 2m_1\tilde{Q}^{bc}Q_{bc} - 2\epsilon^{cc'}\tilde{Q}_{bc}\Phi_2^{bb'}Q_{b'c'} - 2\epsilon^{bb'}\tilde{Q}_{bc}\Phi_3^{cc'}Q_{b'c'} \tag{A.6}$$

of the bifundamental hyper of mass m_1. Here we made use of the identity $\sigma_G^{-1}T\sigma_G = -T^t$.

Remark that the identifications in equation (A.5) reduce the $SU(2)$ R-symmetry for the trifund to that of the bifund, while identifying the gauge symmetry of the first $SU(2)$ gauge group with the enhanced flavor symmetry of the bifund. Also notice that the mass-term breaks the enhanced flavor symmetry of the bifund.

Let us continue by Higgsing the second gauge group by setting $(\Phi_2)^b{}_{b'} = m_2(\sigma_3)^b{}_{b'}$. The single bifundamental hyper turns into two fundamental hypers

$$\begin{aligned} Q_{(k)c} &\equiv Q_{kc} \\ \tilde{Q}_{(k)c} &\equiv \tilde{Q}_{kc}, \end{aligned} \tag{A.7}$$

where a subscript (.) refer to a flavor index. The Yukawa terms can be repackaged as

$$\begin{aligned} W = {} & m_1\,(\sigma_3)^g{}_f\,\delta^l{}_k\,Q^{(f)(k)c}Q_{(g)(l)c} + m_2\,\delta^g{}_f\,(\sigma_3)^l{}_k\,Q^{(f)(k)c}Q_{(g)(l)c} \\ & + \epsilon^{fg}\,\epsilon^{kl}Q_{(f)(k)c}\Phi_3^{cc'}Q_{(g)(l)c'}, \end{aligned} \tag{A.8}$$

if we furthermore make the identifications $Q = Q_{(f=1)}$ and $\tilde{Q} = Q_{(f=2)}$. This superpotential describes two fundamental hypers whose flavor symmetry enhances to $SO(4)$ when the masses are turned off.

As a consistency check let us Higgs both bifundamental gauge groups by setting $(\Phi_2)^b{}_{b'} = m_2(\sigma_3)^b{}_{b'}$ and $(\Phi_3)^c{}_{c'} = m_3(\sigma_3)^c{}_{c'}$ in equation (A.6). This results in the Yukawa terms

$$\begin{aligned} W = {} & m_1\,\delta^l{}_k\,\delta^n{}_m\,(\sigma_3)^g{}_f\,Q^{(f)(k)(m)}Q_{(g)(l)(n)} + m_2\,(\sigma_3)^l{}_k\,\delta^n{}_m\,\delta^g{}_f\,Q^{(f)(k)(m)}Q_{(g)(l)(n)} \\ & + m_3\,\delta^l{}_k\,(\sigma_3)^n{}_m\,\delta^g{}_f\,Q^{(f)(k)(m)}Q_{(g)(l)(n)}, \end{aligned} \tag{A.9}$$

corresponding to the superpotential of eight half-hypers with a diagonal mass matrix with eigenvalues $\pm m_1 \pm m_2 \pm m_3$, as expected.

B. Contour integrands for Sicilian quivers

In this appendix we summarize the contour integrand formulae for the Sicilian quiver gauge theories that we encountered in the main text. More details on instanton counting can be found in [4].

$Sp(1) - SO(4)$ bifundamental full hypermultiplet

The contour integrand for the massive full $Sp(1) - SO(4)$ bifundamental hyper is

$$\begin{aligned}\mathbf{z}^{Sp(1)-SO(4)}_{\text{2bif},\mathrm{k}_1,\mathrm{k}_2}(\phi,\psi,\mu,a,b_1,b_2) = &\left(\prod_{l=1}^{2}\Delta_1(\mu\pm b_l)\right)\Delta_2(\mu\pm a)P_2(\mu)^{\chi_\phi} \\ &\times\left(\frac{\Delta(\mu-\epsilon_-)\Delta(\mu+\epsilon_-)}{\Delta(\mu+\epsilon_+)\Delta(\mu-\epsilon_+)}\right)\left(\frac{\Delta_2(\mu-\epsilon_-)\Delta_2(\mu+\epsilon_-)}{\Delta_2(\mu+\epsilon_+)\Delta_2(\mu-\epsilon_+)}\right)^{\chi_\phi},\end{aligned} \tag{B.1}$$

where the $Sp(1)$ instanton parameter $k_1 = 2n_1 + \chi_\phi$, the deformation parameters $\epsilon_\pm = \frac{\epsilon_1 \pm \epsilon_2}{2}$ and $\pm$ is an abbreviation for a product over both terms. Furthermore, μ is the physical mass parameter and

$$\begin{aligned}\Delta_1(x) &= \prod_{i=1}^{n_1}(\phi_i^2 - x^2) \\ \Delta_2(x) &= \prod_{j=1}^{k_2}(\psi_j^2 - x^2) \\ \Delta(x) &= \prod_{i,j=1}^{n_1,k_2}\left((\phi_i+\psi_j)^2 - x^2\right)\left((\phi_i-\psi_j)^2 - x^2\right) \\ P_2(x,b) &= \prod_{l=1}^{n_2}(b_l^2 - x^2).\end{aligned}$$

$Sp(1) - SO(4)$ single bifundamental half-hypermultiplet

The $Sp(1) - SO(4)$ double bifundamental integrand (B.1) becomes a complete square when the bifundamental mass vanishes. The integrand for the bifundamental half-hyper is thus simply

$$\mathbf{z}^{Sp(1)-SO(4)}_{\text{bif},\mathrm{k}_1,\mathrm{k}_2}(\phi,\psi,a,b_1,b_2) = \prod_{i=1}^{n_1}\left(\phi_i^2 - b_1^2\right)\left(\phi_i^2 - b_2^2\right)\prod_{j=1}^{k_2}\left(a^2 - \psi_j^2\right)\frac{\Delta(\epsilon_-)}{\Delta(\epsilon_+)}\left(b_1 b_2\frac{\Delta_2(\epsilon_-)}{\Delta_2(\epsilon_+)}\right)^{\chi_\phi}.$$

The factor $\Delta(\epsilon_+)$ in the denominator cannot be canceled by a contribution from the gauge multiplets, and therefore brings in additional poles.

$U(2)^3$ trifundamental full hypermultiplet

Starting from the equivariant index

$$\mathrm{Ind}_{\mathbf{T}} = \int_{\mathbb{C}^2} \mathrm{Ch}_{\mathbf{T}}(\mathcal{E}_{U(N_1)} \otimes \mathcal{E}_{U(N_2)} \otimes \mathcal{E}_{U(N_3)} \otimes \mathcal{L} \otimes M)\, \mathrm{Td}_{\mathbf{T}}(\mathbb{C}^2)\,, \tag{B.2}$$

we obtain the contour integrand

$$\begin{aligned}
\mathbf{z}^{(N_1,N_2,N_3)}_{k_1,k_2,k_3} = & \prod_{i,m,n}^{k_1,N_2,N_3} (\phi_{1,i} + b_m + c_n + \mu) \prod_{j,l,n}^{k_2,N_1,N_3} (\phi_{2,j} + a_l + c_n + \mu) \prod_{k,l,m}^{k_3,N_1,N_2} (\phi_{3,k} + a_l + b_m + \mu) \\
& \times \prod_{i,j,n}^{k_1,k_2,N_3} \frac{(\phi_{1,i} + \phi_{2,j} + c_n - \epsilon_- + \mu)(\phi_{1,i} + \phi_{2,j} + c_n + \epsilon_- + \mu)}{(\phi_{1,i} + \phi_{2,j} + c_n + \mu + \epsilon_+)(\phi_{1,i} + \phi_{2,j} + c_n + \mu - \epsilon_+)} \\
& \times \prod_{i,k,m}^{k_2,k_3,N_1} \frac{(\phi_{1,i} + \phi_{3,k} + b_m - \epsilon_- + \mu)(\phi_{1,i} + \phi_{3,k} + b_m + \epsilon_- + \mu)}{(\phi_{1,i} + \phi_{3,k} + b_m + \mu + \epsilon_+)(\phi_{1,i} + \phi_{3,k} + b_m + \mu - \epsilon_+)} \\
& \times \prod_{j,k,l}^{k_2,k_3,N_1} \frac{(\phi_{2,j} + \phi_{3,k} + a_l - \epsilon_- + \mu)(\phi_{2,j} + \phi_{3,k} + a_l + \epsilon_- + \mu)}{(\phi_{2,j} + \phi_{3,k} + a_l + \mu + \epsilon_+)(\phi_{2,j} + \phi_{3,k} + a_l + \mu - \epsilon_+)} \\
& \times \prod_{i,j,k}^{k_1,k_2,k_3} \frac{(\phi^{123}_{ijk} + \epsilon + \mu)(\phi^{123}_{ijk} + \epsilon_1 - \epsilon_2 + \mu)(\phi^{123}_{ijk} - \epsilon_1 + \epsilon_2 + \mu)(\phi^{123}_{ijk} + \mu)^4(\phi^{123}_{ijk} + \mu - \epsilon)}{(\phi^{123}_{ijk} + \epsilon_1 + \mu)^2(\phi^{123}_{ijk} + \epsilon_2 + \mu)^2(\phi^{123}_{ijk} - \epsilon_1 + \mu)^2(\phi^{123}_{ijk} - \epsilon_2 + \mu)^2}
\end{aligned} \tag{B.3}$$

where $\phi^{123}_{ijk} = \phi_{1,i} + \phi_{2,j} + \phi_{3,k}$ and $\epsilon = \epsilon_1 + \epsilon_2$ and $\epsilon_\pm = \frac{\epsilon_1 \pm \epsilon_2}{2}$. The contour integrand for the massive full $SU(2)$ trifundamental hypermultiplet is found by setting $N_{1/2/3} = 2$. If we set the instanton parameter $k_3 = 0$, we recover the contour integrand for two copies of the bifundamental of mass $\pm c$.

The one-loop contribution to the Nekrasov partition function

$$Z^{\text{2trif}}_{\text{1-loop}} = \prod_{i,j=1}^{\infty} \prod_{l,m,n=1}^{2} (a_l + b_m + c_n + \mu + \epsilon_+ + i\epsilon_1 + j\epsilon_2)^{-1}$$

is obtained from the perturbative contribution to the equivariant index (B.2).

References

[1] L. F. Alday, D. Gaiotto and Y. Tachikawa, "Liouville Correlation Functions from Four-Dimensional Gauge Theories," Lett. Math. Phys. **91** (2010) 167 [arXiv:0906.3219 [hep-th]].

[2] D. Gaiotto, "$\mathcal{N}=2$ Dualities," arXiv:0904.2715 [hep-th].

[3] N. Wyllard, "A(N-1) conformal Toda field theory correlation functions from conformal N = 2 SU(N) quiver gauge theories," JHEP **0911**, 002 (2009). [arXiv:0907.2189 [hep-th]].

[4] L. Hollands, C. A. Keller and J. Song, "From SO/Sp Instantons to W-Algebra Blocks," JHEP **1103** (2011) 053 [arXiv:1012.4468 [hep-th]].

[5] O. Chacaltana and J. Distler, "Tinkertoys for Gaiotto Duality," JHEP **1011** (2010) 099 [arXiv:1008.5203 [hep-th]].

[6] P. C. Argyres, N. Seiberg, "S-duality in N=2 supersymmetric gauge theories," JHEP **0712** (2007) 088. [arXiv:0711.0054 [hep-th]].

[7] G. Bonelli, K. Maruyoshi, A. Tanzini, F. Yagi, "Generalized matrix models and AGT correspondence at all genera," [arXiv:1011.5417 [hep-th]].

[8] N. Nekrasov and S. Shadchin, "ABCD of Instantons," Commun. Math. Phys. **252** (2004) 359 [arXiv:hep-th/0404225].

[9] Y. Tachikawa, "Six-dimensional D_N theory and four-dimensional $SO - USp$ quivers," arXiv:0905.4074 [hep-th].

[10] O. Chacaltana, J. Distler, "Tinkertoys for the D_N series," [arXiv:1106.5410 [hep-th]].

[11] D. Gaiotto, "Asymptotically free N=2 theories and irregular conformal blocks," [arXiv:0908.0307 [hep-th]].

[12] E. Witten, "An $SU(2)$ Anomaly," Phys. Lett. B **117** (1982) 324.

[13] N. A. Nekrasov, "Seiberg-Witten Prepotential from Instanton Counting," Adv. Theor. Math. Phys. **7** (2004) 831 [arXiv:hep-th/0206161].

[14] S. Shadchin, "On Certain Aspects of String Theory / Gauge Theory Correspondence," arXiv:hep-th/0502180.

[15] M. R. Gaberdiel, A. Neitzke, "Rationality, quasirationality and finite W-algebras," Commun. Math. Phys. **238** (2003) 305-331. [hep-th/0009235].

[16] A. Marshakov, A. Mironov, A. Morozov, "On non-conformal limit of the AGT relations," Phys. Lett. **B682**, 125-129 (2009). [arXiv:0909.2052 [hep-th]].

[17] S. Cecotti, C. Vafa, "Classification of complete N=2 supersymmetric theories in 4 dimensions," [arXiv:1103.5832 [hep-th]].

[18] U. Bruzzo, F. Fucito, J. F. Morales and A. Tanzini, "Multi-Instanton Calculus and Equivariant Cohomology," JHEP **0305** (2003) 054 [arXiv:hep-th/0211108].

[19] G. W. Moore, N. Nekrasov and S. Shatashvili, "D-Particle Bound States and Generalized Instantons," Commun. Math. Phys. **209** (2000) 77 [arXiv:hep-th/9803265].

[20] M. Gaberdiel, "A general transformation formula for conformal fields," Phys. Lett. **B325** (1994) 366-370. [hep-th/9401166].

[21] F. Benini, S. Benvenuti and Y. Tachikawa, "Webs of five-branes and N=2 superconformal field theories," JHEP **0909**, 052 (2009) [arXiv:0906.0359 [hep-th]].

[22] C. Kozcaz, S. Pasquetti and N. Wyllard, "A & B model approaches to surface operators and Toda theories," JHEP **1008**, 042 (2010) [arXiv:1004.2025 [hep-th]].

[23] N. Nekrasov and V. Pestun, to appear.